arXiv:1609.00705v1 [math.AP] 2 Sep 2016

Classification of the stable solution to the fractional $(2 < s < 3)$ Lane-Emden equation*

Senping Luo[1], **Juncheng Wei**[2] **and Wenming Zou**[3]

[1,3]*Department of Mathematical Sciences, Tsinghua University, Beijing 100084, China*

[2]*Department of Mathematics, University of British Columbia, Vancouver, BC V6T 1Z2, Canada*

Abstract

We classify the stable solutions (positive or sign-changing, radial or not) to the following nonlocal Lane-Emden equation:

$$(-\Delta)^s u = |u|^{p-1}u \quad \text{in} \quad \mathbb{R}^n$$

for $2 < s < 3$.

1 Introduction and Main results

Consider the stable solution of the following equation

$$(-\Delta)^s u = |u|^{p-1}u \quad \text{in} \quad \mathbb{R}^n, \tag{1.1}$$

where $(-\Delta)^s$ is the fractional Laplacian operator for $2 < s < 3$.

The motivation of studying such an equation is originated from the classical Lane-Emden equation

$$-\Delta u = |u|^{p-1}u \quad \text{in} \quad \mathbb{R}^n \tag{1.2}$$

and its parabolic counterpart, which have played a crucial role in the development of nonlinear PDEs in the last decades. These arise in astrophysics and Riemannian geometry. The pioneering works on Eq.(1.2) were contributed by R. Fowler [12, 13]. Later, the ground-breaking result on equation (1.2) is the fundamental Liouville-type theorems established by Gidas and Spruck [14], they claimed that the Eq. (1.2) has no positive solution whenever $p \in (1, 2^* - 1)$, where $2^* = 2n/(n-2)$ if $n \geq 3$ and $2^* = \infty$ if $n \leq 2$. The critical case $p = 2^* - 1$, Eq.(1.2) has a unique positive solution

*Partially supported by NSFC of China and NSERC of Canada. E-mails: luosp14@mails.tsinghua.edu.cn(Luo); jcwei@math.ubc.ca (Wei); wzou@math.tsinghua.edu.cn(Zou)

up to translation and rescaling which is radial and explicitly formulated, see Caffarelli-Gidas-Spruck [1]. Since then many experts in partial differential equations devote to the above equations for various parameters s and p.

For the nonlocal case of $0 < s < 1$, a counterpart of the classification results of Gidas and Spruck [14], and Caffarelli- Gidas-Spruck [1] holds for the fractional Lane-Emden equation (1.1), see the works due to Li [19] and Chen-Li-Ou [5]. In these cases, the Sobolev exponent is given by $P_S(n,s) = (n+2s)/(n-2s)$ if $n > 2s$, and otherwise $P_S(n,s) = \infty$.

Recently, for the nonlocal case of $0 < s < 1$, Davila, Dupaigne and Wei in [6] gave a complete classification of finite Morse index solution of (1.1); for the nonlocal case of $1 < s < 2$, Fazly and Wei in [17] gave a complete classification of finite Morse index solution of (1.1). For the local cases $s = 1$ and $s = 2$, such kind of classification is proved by Farina in [10] and Davila, Dupaigne, Wang and Wei in [7], respectively. For the case $s = 3$, the Joseph-Lundgren exponent (for the triharmonic Lane-Emden equation) is obtained and classification is proved by in [21].

However, when $2 < s < 3$, the equation (1.1) has not been considered so far. In this paper we classify the stable solution of (1.1).

There are many ways of defining the fractional Laplacian $(-\Delta)^s$, where s is any positive, noninteger number. Caffarelli and Silvestre in [2] gave a characterization of the fractional Laplacian when $0 < s < 1$ as the Dirichlet-to-Neumann map for a function u_e satisfying a higher order elliptic equation in the upper half space with one extra spatial dimension. This idea was later generalized by Yang in [27] when the s is being any positive, noninteger number. See also Chang-Gonzales [4] and Case-Chang [3] for general manifolds.

To introduce the fractional operator $(-\Delta)^s$ for $2 < s < 3$, just like the case of $1 < s < 2$, via the Fourier transform, we can define

$$\widehat{(-\Delta)^s u}(\xi) = |\xi|^{2s}\widehat{u}(\xi)$$

or equivalently define this operator inductively by $(-\Delta)^s = (-\Delta)^{s-2} \circ (-\Delta)^2$.

Definition 1.1. *We say a solution u of* (1.1) *is stable outside a compact set if there exists $R_0 > 0$ such that*

$$\int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \frac{(\varphi(x)-\varphi(y))^2}{|x-y|^{n+2s}}dxdy - p\int_{\mathbb{R}^n} |u|^{p-1}\varphi^2 dx \geq 0 \tag{1.3}$$

for any $\varphi \in C_c^\infty(\mathbb{R}^n \backslash \overline{B_{R_0}})$.

Set

$$p_s(n) = \begin{cases} \infty & \text{if } \ n \leq 2s, \\ \frac{n+2s}{n-2s} & \text{if } \ n > 2s. \end{cases}$$

The first main result of the present paper is the following

Theorem 1.1. *Suppose that $n > 2s$ and $2 < s < \delta < 3$. Let $u \in C^{2\delta}(\mathbb{R}^n) \cap L^1(\mathbb{R}^n, (1+|z|)^{n+2s}dz)$ be a solution of* (1.1) *which is stable outside a compact set. Assume*

(1) $1 < p < p_s(n)$ *or*

(2) $p_s(n) < p$ *and*

$$p\frac{\Gamma(\frac{n}{2}-\frac{s}{p-1})\Gamma(s+\frac{s}{p-1})}{\Gamma(\frac{s}{p-1})\Gamma(\frac{n-2s}{2}-\frac{s}{p-1})} > \frac{\Gamma(\frac{n+2s}{4})^2}{\Gamma(\frac{n-2s}{4})^2}, \tag{1.4}$$

then the solution $u \equiv 0$.

(3) $p = p_s(n)$*, then u has finite energy, i.e.,*

$$\|u\|^2_{\dot{H}^s(\mathbb{R}^n)} := \int_{\mathbb{R}^n}\int_{\mathbb{R}^n} \frac{(u(x)-u(y))^2}{|x-y|^{n+2s}}dxdy = \int_{\mathbb{R}^n} |u|^{p+1} < +\infty.$$

If in addition u is stable, then $u \equiv 0$.

Remark 1.1. *In the Theorem 1.1 the condition* (1.4) *is optimal. In fact, the radial singular solution $u = |x|^{-\frac{2s}{p-1}}$ is stable if*

$$p\frac{\Gamma(\frac{n}{2}-\frac{s}{p-1})\Gamma(s+\frac{s}{p-1})}{\Gamma(\frac{s}{p-1})\Gamma(\frac{n-2s}{2}-\frac{s}{p-1})} \le \frac{\Gamma(\frac{n+2s}{4})^2}{\Gamma(\frac{n-2s}{4})^2}.$$

See [22].

Remark 1.2. *The hypothesis* (2) *of Theorem 1.1 is equivalent to*

$$p < p_c(n) := \begin{cases} +\infty & \text{if} \quad n \le n_0(s), \\ \frac{n+2s-2-2a_{n,s}\sqrt{n}}{n-2s-2-2a_{n,s}\sqrt{n}} & \text{if} \quad n > n_0(s), \end{cases} \tag{1.5}$$

where $n_0(s)$ is the largest root of $n-2s-2-2a_{n,s}\sqrt{n}=0$, see [22]. More details and further sharp results about $a_{n,s}$ and $n_0(s)$ see [23].

Remark 1.3. *In this remark, we further analyze the hypothesis* (2) *in Theorem 1.1. Recall that when $s=1$ the condition* (1.4) *gives a upper bounded of p (originated from Joseph and Lundgren [18]), it is*

$$p < p_c(n) := \begin{cases} \infty & \text{if} \quad n \le 10, \\ \frac{(n-2)^2-4n+8\sqrt{n-1}}{(n-2)(n-10)} & \text{if} \quad n \ge 11. \end{cases} \tag{1.6}$$

For the case $s=2$, (1.4) *induce the upper bound of p which is given by the following formula (cf. Gazzola and Grunau [16]):*

$$p < p_c(n) = \begin{cases} \infty & \text{if} \quad n \le 12, \\ \frac{n+2-\sqrt{n^2+4-n\sqrt{n^2-8n+32}}}{n-6-\sqrt{n^2+4-n\sqrt{n^2-8n+32}}} & \text{if} \quad n \ge 13. \end{cases} \tag{1.7}$$

In the triharmonic case, the corresponding exponent given by see ([21]) is the following

$$p < p_c(n) = \begin{cases} \infty & \text{if } n \le 14, \\ \frac{n+4-2D(n)}{n-8-2D(n)} & \text{if } n \ge 15, \end{cases}$$

where

$$D(n) := \frac{1}{6}\Big(9n^2 + 96 - \frac{1536 + 1152n^2}{D_0(n)} - \frac{3}{2}D_0(n)\Big)^{1/2};$$

$$D_0(n) := -(D_1(n) + 36\sqrt{D_2(n)})^{1/3};$$

$$D_1(n) := -94976 + 20736n + 103104n^2 - 10368n^3 + 1296n^5 - 3024n^4 - 108n^6;$$

$$\begin{aligned} D_2(n) :&= 6131712 - 16644096n^2 + 6915840n^4 - 690432n^6 - 3039232n \\ &\quad + 4818944n^3 - 1936384n^5 + 251136n^7 - 30864n^8 - 4320n^9 \\ &\quad + 1800n^{10} - 216n^{11} + 9n^{12}. \end{aligned}$$

2 Preliminary

Throughout this paper we denote $b := 5 - 2s$ and define the operator

$$\Delta_b w := \Delta w + \frac{b}{y} w_y = y^{-b}\mathbf{div}(y^b \nabla w)$$

for a function $w \in W^{3,2}(\mathbb{R}^{n+1}; y^b dxdy)$. We firstly quote the following result.

Theorem 2.1. *(See [27]) Assume $2 < s < 3$. Let $u_e \in W^{3,2}(\mathbb{R}^{n+1}; y^b dxdy)$ satisfy the equation*

$$\Delta_b^3 u_e = 0 \tag{2.1}$$

on the upper half space for $(x, y) \in \mathbb{R}^n \times \mathbb{R}_+$ (where y is the spacial direction) and the boundary conditions:

$$\begin{aligned} &u_e(x, 0) = f(x), \\ &\lim_{y\to 0} y^b \frac{\partial u_e}{\partial y} = 0, \\ &\frac{\partial^2 u_e}{\partial y^2} \mid_{y=0} = \frac{1}{2s}\Delta_x u_e \mid_{y=0}, \\ &\lim_{y\to 0} C_{n,s} y^b \frac{\partial}{\partial y}\Delta_b^2 u_e = (-\Delta)^s f(x), \end{aligned} \tag{2.2}$$

where $f(x)$ is some function defined on $H^s(\mathbb{R}^n)$. Then we have

$$\int_{\mathbb{R}^n} |\xi|^{2s} |\hat{u}(\xi)|^2 d\xi = C_{n,s} \int_{\mathbb{R}^{n+1}_+} y^b |\nabla \Delta_b u_e(x, y)|^2 dxdy. \tag{2.3}$$

Applying the above theorem to solutions of (1.1), we conclude that the extended function $u_e(x,y)$, where $x=(x_1,...,x_n)\in\mathbb{R}^n$ and $y\in\mathbb{R}^+$, satisfies

$$\begin{cases}\Delta_b^3 u_e=0 \ \text{ in } \ \mathbb{R}_+^{n+1},\\ \lim_{y\to 0} y^b\frac{\partial u_e}{\partial y}=0 \ \text{ on } \ \partial\mathbb{R}_+^{n+1},\\ \frac{\partial^2 u_e}{\partial y^2}\mid_{y=0}=\frac{1}{2s}\Delta_x u_e\mid_{y=0} \ \text{ on } \ \partial\mathbb{R}_+^{n+1},\\ \lim_{y\to 0} y^b\frac{\partial}{\partial y}\Delta_b^2 u_e=-C_{n,s}|u_e|^{p-1}u_e \ \text{ in } \ \mathbb{R}_+^{n+1}.\end{cases} \tag{2.4}$$

Moreover,

$$\int_{\mathbb{R}^n}|\xi|^{2s}|\hat{u}(\xi)|^2d\xi=C_{n,s}\int_{\mathbb{R}_+^{n+1}}y^b|\nabla\Delta_b u_e(x,y)|^2dxdy$$

and $u(x)=u_e(x,0)$.

Define

$$\begin{aligned}E(\lambda,x,u_e)=&\int_{\mathbb{R}_+^{n+1}\cap\partial B_1}\frac{1}{2}\theta_1^b|\nabla\Delta_b u_e^\lambda|^2-\frac{C_{n,s}}{p+1}\int_{\partial\mathbb{R}_+^{n+1}\cap B_1}|u_e^\lambda|^{p+1}\\ &+\sum_{0\le i,j\le 4,i+j\le 5}C_{i,j}^1\int_{\mathbb{R}_+^{n+1}\cap\partial B_1}\theta_1^b\lambda^{i+j}\frac{d^iu_e^\lambda}{d\lambda^i}\frac{d^ju_e^\lambda}{d\lambda^j}\\ &+\sum_{0\le t,s\le 2,t+s\le 3}C_{t,s}^2\int_{\mathbb{R}_+^{n+1}\cap\partial B_1}\theta_1^b\lambda^{t+s}\nabla_{S^n}\frac{d^tu_e^\lambda}{d\lambda^t}\nabla_{S^n}\frac{d^su_e^\lambda}{d\lambda^s}\\ &+\sum_{0\le l,k\le 1,l+k\le 1}C_{l,k}^3\int_{\mathbb{R}_+^{n+1}\cap\partial B_1}\theta_1^b\lambda^{l+k}\Delta_{S^n}\frac{d^lu_e^\lambda}{d\lambda^l}\Delta_{S^n}\frac{d^ku_e^\lambda}{d\lambda^k}\\ &+(\frac{s}{p-1}+1)\int_{\mathbb{R}_+^{n+1}\cap\partial B_1}\theta_1^b(\Delta_b u_e^\lambda)^2.\end{aligned} \tag{2.5}$$

The following is the monotonicity formula which will paly an important role.

Theorem 2.2. *Let u_e satisfy the equation* (2.1) *with the boundary conditions* (2.2)*, we have the following*

$$\begin{aligned}\frac{dE(\lambda,x,u_e)}{d\lambda}=&\int_{\mathbb{R}_+^{n+1}\cap\partial B_1}\theta_1^b\Big(3\lambda^5(\frac{d^3u_e^\lambda}{d\lambda^3})^2+A_1\lambda^3(\frac{d^2u_e^\lambda}{d\lambda^2})^2+A_2\lambda(\frac{du_e^\lambda}{d\lambda})^2\Big)\\ &+\int_{\mathbb{R}_+^{n+1}\cap\partial B_1}\theta_1^b\Big(2\lambda^3|\nabla_{S^n}\frac{d^2u_e^\lambda}{d\lambda^2}|^2+B_1\lambda|\nabla_{S^n}\frac{du_e^\lambda}{d\lambda}|\Big)\\ &+\int_{\mathbb{R}_+^{n+1}\cap\partial B_1}\theta_1^b\lambda(\Delta_{S^n}\frac{du_e^\lambda}{d\lambda})^2,\end{aligned} \tag{2.6}$$

where $\theta_1 = \frac{y}{r}$ and

$$
\begin{aligned}
&A_1 := 10\delta_1 - 2\delta_2 - 56 + \alpha_0^2 - 2\alpha_0 - 2\beta_0 - 4,\\
&A_2 := -18\delta_1 + 6\delta_2 - 4\delta_3 + 2\delta_4 + 72 - \alpha_0^2 + \beta_0^2 + 2\alpha_0 + 2\beta_0,\\
&B_1 := 8\alpha - 4\beta - 2\beta_0 + 4(n+b) - 14,\\
&\alpha := n + b - 2 - \frac{4s}{p-1}, \beta := \frac{2s}{p-1}(3 + \frac{2s}{p-1} - n - b),\\
&\alpha_0 := n + b - \frac{4s}{p-1}, \quad \beta_0 := \frac{2s}{p-1}(1 + \frac{2s}{p-1} - n - b)
\end{aligned}
$$

and

$$
\begin{aligned}
\delta_1 =& 2(n+b) - \frac{8s}{p-1},\\
\delta_2 =& (n+b)(n+b-2) - (n+b)\frac{12s}{p-1} + \frac{12s}{p-1}(1 + \frac{2s}{p-1}),\\
\delta_3 =& -\frac{8s}{p-1}(1 + \frac{2s}{p-1})(2 + \frac{2s}{p-1}) + 2(n+b)\frac{6s}{p-1}(1 + \frac{2s}{p-1})\\
&- (n+b)(n+b-2)(1 + \frac{4s}{p-1}),\\
\delta_4 =& (3 + \frac{2s}{p-1})(2 + \frac{2s}{p-1})(1 + \frac{2s}{p-1})\frac{2s}{p-1}\\
&- 2(n+b)(1 + \frac{2s}{p-1})(2 + \frac{2s}{p-1})\frac{2s}{p-1}\\
&+ (n+b)(n+b-2)(2 + \frac{2s}{p-1})\frac{2s}{p-1}.
\end{aligned}
\tag{2.7}
$$

We will give the proof of Theorem 2.2 in the next section. Now we would like to state a consequent result of Theorem 2.2. Recall that $E(\lambda, x, u_e)$, defined in (2.5), can be divided into two parts: the integral over the ball B_λ and the terms on the boundary ∂B_λ. We note that in our blow-down analysis, the coefficients (including positive or negative, big or small) of the boundary terms can be estimated in a unified way, therefore we may change some coefficients of the boundary terms in $E(\lambda, x, u_e)$. After such a change, we denote the new functional by $E^c(\lambda, x, u_e)$.

Define

$$
p_m(n) := \begin{cases} +\infty & \text{if} \quad n < 2s + 6 + \sqrt{73},\\ \frac{5n+10s-\sqrt{15(n-2s)^2+120(n-2s)+370}}{5n-10s-\sqrt{15(n-2s)^2+120(n-2s)+370}} & \text{if} \quad n \geq 2s + 6 + \sqrt{73}. \end{cases} \tag{2.8}
$$

We have the following

Theorem 2.3. *Assume that $\frac{n+2s}{n-2s} < p < p_m(n)$, then $E^c(\lambda, x, u_e)$ is a nondecreasing function of $\lambda > 0$. Furthermore,*

$$
\frac{dE^c(\lambda, x, u_e)}{d\lambda} \geq C(n, s, p)\lambda^{2s\frac{p+1}{p-1}-6-n} \int_{\mathbb{R}^{n+1}_+ \cap \partial B_\lambda(x_0)} y^b \Big(\frac{2s}{p-1}u_e + \lambda \partial_r u_e\Big)^2,
$$

where $C(n,s,p)$ is a constant independent of λ.

By carefully comparing $\frac{n+2s}{n-2s} < p < p_m(n)$ with $p > \frac{n+2s}{n-2s}$ and (1.4), we get the following (see the last section of the current paper) monotonicity formula for our blow down analysis.

Theorem 2.4. *Assume that $p > \frac{n+2s}{n-2s}$ and* (1.4)*, then $E^c(\lambda, x, u_e)$ is a nondecreasing function of $\lambda > 0$. Furthermore,*

$$\frac{dE^c(\lambda,x,u_e)}{d\lambda} \geq C(n,s,p)\lambda^{2s\frac{p+1}{p-1}-6-n}\int_{\mathbb{R}^{n+1}_+\cap\partial B_\lambda(x_0)} y^b\Big(\frac{2s}{p-1}u_e + \lambda\partial_r u_e\Big)^2,$$

where $C(n,s,p)$ is a constant independent of λ.

3 Monotonicity formula and the proof of Theorem 2.2

The derivation of the monotonicity for the (1.1) when $2 < s < 3$ is complicated in its process, we divide it into several subsections. In subsection 3.1, we derive $\frac{d}{d\lambda}\overline{E}(u_e,\lambda)$. In subsection 3.2, we calculate $\frac{\partial^j}{\partial r^j}u_e^\lambda$ and $\frac{\partial^i}{\partial\lambda^i}u_e^\lambda$, $i,j = 1,2,3,4$. In subsection 3.3, the operator Δ_b^2 and its representation will be given. In subsection 3.4, we decompose $\frac{d}{d\lambda}\overline{E}(u_e^\lambda,1)$. Finally, combine with the above four subsections, we can obtain the monotonicity formula, hence get the proof of Theorem 2.2.

Suppose that $x_0 = 0$ and denote by B_λ the balls centered at zero with radius λ. Set

$$\overline{E}(u_e,\lambda) := \lambda^{2s\frac{p+1}{p-1}-n}\Big(\int_{\mathbb{R}^{n+1}_+\cap B_\lambda}\frac{1}{2}y^b|\nabla\Delta_b u_e|^2 - \frac{C_{n,s}}{p+1}\int_{\partial\mathbb{R}^{n+1}_+\cap B_\lambda}|u_e|^{p+1}\Big).$$

3.1 The derivation of $\frac{d}{d\lambda}\overline{E}(u_e,\lambda)$

Define

$$\begin{aligned} &v_e := \Delta_b u_e, u_e^\lambda(X) := \lambda^{\frac{2s}{p-1}}u_e(\lambda X), \quad w_e(X) := \Delta_b v_e \\ &v_e^\lambda(X) := \lambda^{\frac{2s}{p-1}+2}v_e(\lambda X), \quad w_e^\lambda(X) := \lambda^{\frac{2s}{p-1}+4}w_e(\lambda X), \end{aligned} \tag{3.1}$$

where $X = (x,y) \in \mathbb{R}^{n+1}_+$. Therefore,

$$\Delta_b u_e^\lambda(X) = v_e^\lambda(X), \Delta_b v_e^\lambda(X) = w_e^\lambda(X). \tag{3.2}$$

Hence

$$\begin{aligned} &\Delta_b w_e^\lambda = 0, \\ &\lim_{y\to 0} y^b\frac{\partial u_e}{\partial y} = 0, \\ &\frac{\partial^2 u_e}{\partial y^2}\mid_{y=0} = \frac{1}{2s}\Delta_x u_e\mid_{y=0}, \\ &\lim_{y\to 0} C_{n,s}y^b\frac{\partial}{\partial y}w_e^\lambda = -C_{n,s}|u_e|^{p-1}u_e. \end{aligned}$$

In addition, differentiating (3.2) with respect to λ we have

$$\Delta_b \frac{du_e^\lambda}{d\lambda} = \frac{dv_e^\lambda}{d\lambda}, \quad \Delta_b \frac{dv_e^\lambda}{d\lambda} = \frac{dw_e^\lambda}{d\lambda}.$$

Note that

$$\overline{E}(u_e, \lambda) = \overline{E}(u_e^\lambda, 1) = \int_{\mathbb{R}_+^{n+1}\cap B_1} \frac{1}{2} y^b |\nabla v_e^\lambda|^2 - \frac{C_{n,s}}{p+1} \int_{\partial\mathbb{R}_+^{n+1}\cap B_1} |u_e^\lambda|^{p+1}.$$

Taking derivative of the energy $\overline{E}(u_e^\lambda, 1)$ with respect to λ and integrating by part we have:

$$\begin{aligned}
&\frac{d\overline{E}(u_e^\lambda, 1)}{d\lambda} = \int_{\mathbb{R}_+^{n+1}\cap B_1} y^b \nabla v_e^\lambda \nabla \frac{dv_e^\lambda}{d\lambda} - C_{n,s} \int_{\partial\mathbb{R}_+^{n+1}\cap B_1} |u_e^\lambda|^{p-1} u_e^\lambda \frac{du_e^\lambda}{d\lambda} \\
&= \int_{\partial(\mathbb{R}_+^{n+1}\cap B_1)} y^b \frac{\partial v_e^\lambda}{\partial n} \frac{dv_e^\lambda}{d\lambda} - \int_{\mathbb{R}_+^{n+1}\cap B_1} (y^b \Delta v_e^\lambda + b y^{b-1} \frac{\partial v_e^\lambda}{\partial y}) \frac{dv_e^\lambda}{d\lambda} \\
&\quad - C_{n,s} \int_{\partial\mathbb{R}_+^{n+1}\cap B_1} |u_e^\lambda|^{p-1} u_e^\lambda \frac{du_e^\lambda}{d\lambda} \qquad (3.3) \\
&= - \int_{\partial\mathbb{R}^{n+1}\cap B_1} y^b \frac{\partial v_e^\lambda}{\partial y} \frac{dv_e^\lambda}{d\lambda} + \int_{\mathbb{R}_+^{n+1}\cap \partial B_1} y^b \frac{\partial v_e^\lambda}{\partial r} \\
&\quad - \int_{\mathbb{R}_+^{n+1}\cap B_1} y^b \Delta v_e^\lambda \frac{dv_e^\lambda}{d\lambda} + b y^{b-1} \frac{\partial v_e^\lambda}{\partial y} \frac{v_e^\lambda}{d\lambda} - \int_{\partial\mathbb{R}_+^{n+1}\cap B_1} y^b \frac{w_e^\lambda}{\partial y} \frac{u_e^\lambda}{d\lambda}.
\end{aligned}$$

Now note that from the definition of v_e^λ and by differentiating it with respect to λ, we get the following identity for $X \in \mathbb{R}_+^{n+1}$,

$$r \frac{\partial v_e^\lambda}{\partial r} = \lambda \partial_\lambda v_e^\lambda - (\frac{2s}{p-1} + 2) v_e^\lambda.$$

Hence,

$$\begin{aligned}
\int_{\mathbb{R}_+^{n+1}\cap B_1} y^b \frac{\partial v_e^\lambda}{\partial r} \frac{dv_e^\lambda}{d\lambda} &= \int_{\mathbb{R}_+^{n+1}\cap B_1} y^b \Big(\lambda \frac{dv_e^\lambda}{d\lambda} \frac{dv_e^\lambda}{d\lambda} - (\frac{2s}{p-1} + 2) v_e^\lambda \frac{dv_e^\lambda}{d\lambda} \Big) \\
&= \lambda \int_{\mathbb{R}_+^{n+1}\cap \partial B_1} y^b (\frac{dv_e^\lambda}{d\lambda})^2 - (\frac{s}{p-1} + 1) \frac{d}{d\lambda} \int_{\mathbb{R}_+^{n+1}\cap \partial B_1} y^b (v_e^\lambda)^2.
\end{aligned}$$

Note that

$$\begin{aligned}
- \int_{\mathbb{R}_+^{n+1}\cap B_1} y^b \Delta v_e^\lambda \frac{dv_e^\lambda}{d\lambda} &= \int_{\partial\mathbb{R}_+^{n+1}\cap B_1} y^b \frac{\partial v_e^\lambda}{\partial y} \frac{v_e^\lambda}{d\lambda} - \int_{\mathbb{R}_+^{n+1}\cap \partial B_1} y^b \frac{\partial v_e^\lambda}{\partial r} \frac{v_e^\lambda}{d\lambda} \\
&\quad + \int_{\mathbb{R}_+^{n+1}\cap B_1} \nabla v_e^\lambda \nabla (y^b \frac{dv_e^\lambda}{d\lambda}).
\end{aligned}$$

Integration by part we have

$$\begin{aligned}
&\int_{\mathbb{R}^{n+1}_+\cap B_1} y^b \nabla v_e^\lambda \nabla \frac{dv_e^\lambda}{d\lambda} \\
&= -\int_{\partial\mathbb{R}^{n+1}_+\cap B_1} y^b \frac{\partial v_e^\lambda}{\partial y}\frac{dv_e^\lambda}{d\lambda} + \int_{\mathbb{R}^{n+1}_+\cap \partial B_1} y^b \frac{\partial v_e^\lambda}{\partial r}\frac{dv_e^\lambda}{d\lambda} \\
&\quad -\int_{\mathbb{R}^{n+1}_+\cap B_1} \nabla\cdot(y^b\nabla v_e^\lambda)\frac{dv_e^\lambda}{d\lambda} \\
&= -\int_{\partial\mathbb{R}^{n+1}_+\cap B_1} y^b \frac{\partial v_e^\lambda}{\partial y}\frac{dv_e^\lambda}{d\lambda} + \int_{\mathbb{R}^{n+1}_+\cap \partial B_1} y^b \frac{\partial v_e^\lambda}{\partial r}\frac{dv_e^\lambda}{d\lambda} \\
&\quad -\int_{\mathbb{R}^{n+1}_+\cap B_1} y^b \Delta_b v_e^\lambda \Delta_b \frac{du_e^\lambda}{d\lambda}.
\end{aligned}$$

Now

$$\begin{aligned}
-\int_{\mathbb{R}^{n+1}_+\cap B_1} y^b \Delta_b v_e^\lambda \Delta_b \frac{du_e^\lambda}{d\lambda} &= -\int_{\mathbb{R}^{n+1}_+\cap B_1} y^b \Delta_b v_e^\lambda (\Delta \frac{du_e\lambda}{d\lambda} + \frac{b}{y}\frac{\partial}{\partial y}\frac{du_e^\lambda}{d\lambda}) \\
&= -\int_{\partial(\mathbb{R}^{n+1}_+\cap B_1)} y^b \Delta_b v_e^\lambda \frac{\partial}{\partial n}\frac{du_e^\lambda}{d\lambda} + \int_{\mathbb{R}^{n+1}_+\cap B_1} \nabla(y^b\Delta_b v_e^\lambda)\nabla\frac{du_e^\lambda}{d\lambda} \\
&\quad -\int_{\mathbb{R}^{n+1}_+\cap B_1} b y^{b-1}\Delta_b v_e^\lambda \frac{\partial}{\partial y}\frac{du_e^\lambda}{d\lambda} \\
&= -\int_{\partial(\mathbb{R}^{n+1}_+\cap B_1)} y^b \Delta_b v_e^\lambda \frac{\partial}{\partial n}\frac{du_e^\lambda}{d\lambda} + \int_{\mathbb{R}^{n+1}_+\cap B_1} y^b \nabla\Delta_b v_e^\lambda \nabla\frac{du_e^\lambda}{d\lambda} \\
&= -\int_{\partial(\mathbb{R}^{n+1}_+\cap B_1)} y^b \Delta_b v_e^\lambda \frac{\partial}{\partial n}\frac{du_e^\lambda}{d\lambda} + \int_{\partial(\mathbb{R}^{n+1}_+\cap B_1)} y^b \frac{\partial\Delta_b v_e^\lambda}{\partial n}\frac{du_e^\lambda}{d\lambda} \\
&\quad -\int_{\mathbb{R}^{n+1}_+\cap B_1} y^b \Delta_b^2 v_e^\lambda \frac{du_e^\lambda}{d\lambda} \\
&= -\int_{\partial(\mathbb{R}^{n+1}_+\cap B_1)} y^b \Delta_b v_e^\lambda \frac{\partial}{\partial n}\frac{du_e^\lambda}{d\lambda} + \int_{\partial(\mathbb{R}^{n+1}_+\cap B_1)} y^b \frac{\partial\Delta_b v_e^\lambda}{\partial n}\frac{du_e^\lambda}{d\lambda}.
\end{aligned}$$

Here we have used that $\Delta_b^2 v_e^\lambda = \Delta_b^3 u_e^\lambda = 0$. Therefore, combine with the above arguments we get that

$$\begin{aligned}\int_{\mathbb{R}_+^{n+1}\cap B_1} y^b \nabla v_e^\lambda \nabla \frac{dv_e^\lambda}{d\lambda} &= -\int_{\partial\mathbb{R}_+^{n+1}\cap B_1} y^b \frac{\partial v_e^\lambda}{\partial y}\frac{dv_e^\lambda}{d\lambda} + \int_{\mathbb{R}_+^{n+1}\cap \partial B_1} y^b \frac{\partial v_e^\lambda}{\partial r}\frac{dv_e^\lambda}{d\lambda}\\ &+ \int_{\partial\mathbb{R}_+^{n+1}\cap B_1} y^b \Delta_b v_e^\lambda \frac{\partial}{\partial y}\frac{du_e^\lambda}{d\lambda} - \int_{\mathbb{R}_+^{n+1}\cap \partial B_1} y^b \Delta v_e^\lambda \frac{\partial}{\partial r}\frac{du_e^\lambda}{d\lambda}\\ &- \int_{\partial\mathbb{R}_+^{n+1}\cap B_1} y^b \frac{\partial}{\partial y}\Delta_b v_e^\lambda \frac{du_e^\lambda}{d\lambda} + \int_{\mathbb{R}_+^{n+1}\cap \partial B_1} y^b \frac{\partial}{\partial r}\Delta_b v_e^\lambda \frac{du_e^\lambda}{d\lambda}\\ &= \int_{\mathbb{R}_+^{n+1}\cap \partial B_1} y^b \frac{\partial v_e^\lambda}{\partial r}\frac{dv_e^\lambda}{d\lambda} - \int_{\mathbb{R}_+^{n+1}\cap \partial B_1} y^b \Delta v_e^\lambda \frac{\partial}{\partial r}\frac{du_e^\lambda}{d\lambda}\\ &- C(n,s)\int_{\partial\mathbb{R}_+^{n+1}\cap B_1} |u_e^\lambda|^{p-1}\frac{du_e^\lambda}{d\lambda} + \int_{\mathbb{R}_+^{n+1}\cap \partial B_1} y^b \frac{\partial}{\partial r}\Delta_b v_e^\lambda \frac{du_e^\lambda}{d\lambda}.\end{aligned} \tag{3.4}$$

Here, we have used that $\frac{\partial \Delta u_e^\lambda(x,0)}{\partial y} = 0$, $\frac{\partial}{\partial y}\frac{du_e^\lambda}{d\lambda} = 0$ on $\partial\mathbb{R}_+^{n+1}$ and $\lim_{y\to 0} y^b \frac{\partial}{\partial y}\Delta_b v_e^\lambda = -C_{n,s}|u_e^\lambda|^{p-1}u_e^\lambda$. By (3.3) and (3.4) we obtain that

$$\begin{aligned}\frac{d}{d\lambda}\overline{E}(u_e^\lambda, 1) = &\int_{\mathbb{R}_+^{n+1}\cap \partial B_1} y^b \frac{\partial v_e^\lambda}{\partial r}\frac{dv_e^\lambda}{d\lambda} + \int_{\mathbb{R}_+^{n+1}\cap \partial B_1} y^b \frac{\partial w_e^\lambda}{\partial r}\frac{du_e^\lambda}{d\lambda}\\ &- \int_{\mathbb{R}_+^{n+1}\cap \partial B_1} y^b w_e^\lambda \frac{\partial}{\partial r}\frac{du_e^\lambda}{d\lambda}.\end{aligned} \tag{3.5}$$

Recall (3.1) and differentiate it with respect to λ, we have

$$\begin{aligned}\frac{du_e^\lambda(X)}{d\lambda} &= \frac{1}{\lambda}\big(\frac{2s}{p-1}u_e^\lambda(X) + r\partial_r u_e^\lambda(X)\big),\\ \frac{dv_e^\lambda(X)}{d\lambda} &= \frac{1}{\lambda}\big((\frac{2s}{p-1}+2)v_e^\lambda(X) + r\partial_r v_e^\lambda(X)\big),\\ \frac{dw_e^\lambda(X)}{d\lambda} &= \frac{1}{\lambda}\big((\frac{2s}{p-1}+4)w_e^\lambda(X) + r\partial_r w_e^\lambda(X)\big).\end{aligned}$$

Differentiate the above equations with respect to λ again we get

$$\lambda\frac{d^2u_e^\lambda(X)}{d\lambda^2} + \frac{du_e^\lambda(x)}{d\lambda} = \frac{2s}{p-1}\frac{du_e^\lambda(X)}{d\lambda} + r\partial_r\frac{du_e^\lambda}{d\lambda}.$$

Hence, for $X \in \mathbb{R}^{n+1}_+ \cap B_1$, we have

$$\begin{aligned}
\partial_r(u_e^\lambda(X)) &= \lambda\frac{du_e^\lambda}{d\lambda} - \frac{2s}{p-1}u_e,\\
\partial_r(\frac{du_e^\lambda(X)}{d\lambda}) &= \lambda\frac{d^2u_e^\lambda(X)}{d\lambda^2} + (1-\frac{2s}{p-1})\frac{du_e^\lambda}{d\lambda},\\
\partial_r(v_e^\lambda(X)) &= \lambda\frac{dv_e^\lambda}{d\lambda} - (\frac{2s}{p-1}+2)v_e^\lambda,\\
\partial_r(w_e^\lambda(X)) &= \lambda\frac{dw_e^\lambda}{d\lambda} - (\frac{2s}{p-1}+4)w_e^\lambda.
\end{aligned}$$

Plugging these equations into (3.5), we get that

$$\begin{aligned}
\frac{d}{d\lambda}\overline{E}(u_e^\lambda,1) =& \int_{\mathbb{R}^{n+1}_+\cap\partial B_1} y^b\big(\lambda\frac{dv_e^\lambda}{d\lambda}\frac{dv_e^\lambda}{d\lambda} - (\frac{2s}{p-1}+2)v_e^\lambda\frac{dv_e^\lambda}{d\lambda}\big)\\
&+ y^b\big(\lambda\frac{dw_e^\lambda}{d\lambda}\frac{u_e^\lambda}{d\lambda} - (\frac{2s}{p-1}+4)w_e^\lambda\frac{du_e^\lambda}{d\lambda}\big)\\
&- y^b\big(\lambda w_e^\lambda\frac{d^2u_e^\lambda}{d\lambda^2} + (1-\frac{2s}{p-1}w_e^\lambda\frac{du_e^\lambda}{d\lambda}\big)\\
=& \underbrace{\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} y^b\big[\lambda\frac{dv_e^\lambda}{d\lambda}\frac{dv_e^\lambda}{d\lambda} - (\frac{2s}{p-1}+2)v_e^\lambda\frac{dv_e^\lambda}{d\lambda}\big]}\\
&+ \underbrace{y^b\big[\lambda\frac{dw_e^\lambda}{d\lambda}\frac{du_e^\lambda}{d\lambda} - \lambda w_e^\lambda\frac{d^2u_e^\lambda}{d\lambda^2}\big] - 5y^bw_e^\lambda\frac{du_e^\lambda}{d\lambda}}\\
:=& \overline{E}_{d1}(u_e^\lambda,1) + \overline{E}_{d2}(u_e^\lambda,2).
\end{aligned} \tag{3.6}$$

3.2 The calculations of $\frac{\partial^j}{\partial r^j}u_e^\lambda$ and $\frac{\partial^i}{\partial\lambda^i}u_e^\lambda,\ i,j=1,2,3,4$

Note

$$\lambda\frac{du_e^\lambda}{d\lambda} = \frac{2s}{p-1}u_e^\lambda + r\frac{\partial}{\partial r}u_e^\lambda. \tag{3.7}$$

Differentiating (3.7) once, twice and thrice with respect to λ respectively, we have

$$\lambda\frac{d^2u_e^\lambda}{d\lambda^2} + \frac{du_e^\lambda}{d\lambda} = \frac{2s}{p-1}\frac{du_e^\lambda}{d\lambda} + r\frac{\partial}{\partial r}\frac{du_e^\lambda}{d\lambda}, \tag{3.8}$$

$$\lambda\frac{d^3u_e^\lambda}{d\lambda^3} + 2\frac{d^2u_e^\lambda}{d\lambda^2} = \frac{2s}{p-1}\frac{d^2u_e^\lambda}{d\lambda^2} + r\frac{\partial}{\partial r}\frac{d^2u_e^\lambda}{d\lambda^2}, \tag{3.9}$$

$$\lambda\frac{d^4u_e^\lambda}{d\lambda^4} + 3\frac{d^3u_e^\lambda}{d\lambda^3} = \frac{2s}{p-1}\frac{d^3u_e^\lambda}{d\lambda^3} + r\frac{\partial}{\partial r}\frac{d^3u_e^\lambda}{d\lambda^3}. \tag{3.10}$$

Similarly, differentiating (3.7) once, twice and thrice with respect to r respectively we have

$$\lambda\frac{\partial}{\partial r}\frac{du_e^\lambda}{d\lambda} = (\frac{2s}{p-1}+1)\frac{\partial}{\partial r}u_e^\lambda + r\frac{\partial^2}{\partial r^2}u_e^\lambda, \tag{3.11}$$

$$\lambda\frac{\partial^2}{\partial r^2}\frac{du_e^\lambda}{d\lambda}=(\frac{2s}{p-1}+2)\frac{\partial^2}{\partial r^2}u_e^\lambda+r\frac{\partial^3}{\partial r^3}u_e^\lambda, \tag{3.12}$$

$$\lambda\frac{\partial^3}{\partial r^3}\frac{du_e^\lambda}{d\lambda}=(\frac{2s}{p-1}+3)\frac{\partial^3}{\partial r^3}u_e^\lambda+r\frac{\partial^4}{\partial r^4}u_e^\lambda. \tag{3.13}$$

From (3.7), on $\mathbb{R}_+^{n+1}\cap\partial B_1$, we have

$$\frac{\partial u_e^\lambda}{\partial r}=\lambda\frac{du_e^\lambda}{d\lambda}-\frac{2s}{p-1}u_e^\lambda.$$

Next from (3.8), on $\mathbb{R}_+^{n+1}\cap\partial B_1$, we derive that

$$\frac{\partial}{\partial r}\frac{du_e^\lambda}{d\lambda}=\lambda\frac{d^2u_e^\lambda}{d\lambda^2}+(1-\frac{2s}{p-1})\frac{du_e^\lambda}{d\lambda}.$$

From (3.11), combine with the two equations above, on $\mathbb{R}_+^{n+1}\cap\partial B_1$, we get

$$\begin{aligned}\frac{\partial^2}{\partial r^2}u_e^\lambda&=\lambda\frac{\partial}{\partial r}\frac{du_e^\lambda}{d\lambda}-(1+\frac{2s}{p-1})\frac{\partial}{\partial r}u_e^\lambda\\&=\lambda^2\frac{d^2u_e^\lambda}{d\lambda^2}-\lambda\frac{4s}{p-1}\frac{du_e^\lambda}{d\lambda}+(1+\frac{2s}{p-1})\frac{2s}{p-1}u_e^\lambda.\end{aligned} \tag{3.14}$$

Differentiating (3.8) with respect to r, and combine with (3.8) and (3.9), we get that

$$\begin{aligned}\frac{\partial^2}{\partial r^2}\frac{du_e^\lambda}{d\lambda}&=\lambda\frac{\partial}{\partial r}\frac{d^2u_e^\lambda}{d\lambda^2}-\frac{2s}{p-1}\frac{\partial}{\partial r}\frac{du_e^\lambda}{d\lambda}\\&=\lambda^2\frac{d^3u_e^\lambda}{d\lambda^3}+(2-\frac{4s}{p-1})\lambda\frac{d^2u_e^\lambda}{d\lambda^2}-(1-\frac{2s}{p-1})\frac{2s}{p-1}\frac{du_e^\lambda}{d\lambda}.\end{aligned} \tag{3.15}$$

From (3.12), on $\mathbb{R}_+^{n+1}\cap\partial B_1$, combine with (3.14) and (3.15), we have

$$\begin{aligned}\frac{\partial^3}{\partial r^3}u_e^\lambda=&\lambda\frac{\partial^2}{\partial r^2}\frac{du_e^\lambda}{d\lambda}-(2+\frac{2s}{p-1})\frac{\partial^2}{\partial r^2}u_e^\lambda\\=&\lambda^3\frac{d^3u_e^\lambda}{d\lambda^3}-\lambda^2\frac{6s}{p-1}\frac{d^2u_e^\lambda}{d\lambda^2}+\lambda(\frac{6s}{p-1}+\frac{12s^2}{(p-1)^2})\frac{du_e^\lambda}{d\lambda}\\&-(2+\frac{2s}{p-1})(1+\frac{2s}{p-1})\frac{2s}{p-1}u_e^\lambda.\end{aligned} \tag{3.16}$$

Now differentiating (3.8) once with respect to r, we get

$$\lambda\frac{\partial^2}{\partial r^2}\frac{d^2u_e^\lambda}{d\lambda^2}=(\frac{2s}{p-1}+1)\frac{\partial^2}{\partial r^2}\frac{du_e^\lambda}{d\lambda}+r\frac{\partial^3}{\partial r^3}\frac{du_e^\lambda}{d\lambda},$$

then on $\mathbb{R}_+^{n+1}\cap\partial B_1$, we have

$$\frac{\partial^3}{\partial r^3}\frac{du_e^\lambda}{d\lambda}=\lambda\frac{\partial^2}{\partial r^2}\frac{d^2u_e^\lambda}{d\lambda^2}-(\frac{2s}{p-1}+1)\frac{\partial^2}{\partial r^2}\frac{du_e^\lambda}{d\lambda}. \tag{3.17}$$

Now differentiating (3.9) twice with respect to r, we get

$$\lambda\frac{\partial}{\partial r}\frac{d^3u_e^\lambda}{d\lambda^3}=(\frac{2s}{p-1}-1)\frac{\partial}{\partial r}\frac{d^2u_e^\lambda}{d\lambda^2}+r\frac{\partial^2}{\partial r^2}\frac{d^2u_e^\lambda}{d\lambda^2},$$

hence on $\mathbb{R}_+^{n+1}\cap\partial B_1$, combine with (3.9) and (3.10) there holds

$$\begin{aligned}\frac{\partial^2}{\partial r^2}\frac{d^2u_e^\lambda}{d\lambda^2}=&\lambda\frac{\partial}{\partial r}\frac{d^3u_e^\lambda}{d\lambda^3}+(1-\frac{2s}{p-1})\frac{\partial}{\partial r}\frac{d^2u_e^\lambda}{d\lambda^2}\\ =&\lambda^2\frac{d^4u_e^\lambda}{d\lambda^4}+\lambda(4-\frac{4s}{p-1})\frac{d^3u_e^\lambda}{d\lambda^3}+(1-\frac{2s}{p-1})(2-\frac{2s}{p-1})\frac{d^2u_e^\lambda}{d\lambda^2}.\end{aligned}\tag{3.18}$$

Now differentiating (3.8) with respect to r, we have

$$\lambda\frac{\partial}{\partial r}\frac{d^2u_e^\lambda}{d\lambda^2}=\frac{2s}{p-1}\frac{\partial}{\partial r}\frac{du_e^\lambda}{d\lambda}+r\frac{\partial^2}{\partial r^2}\frac{du_e^\lambda}{d\lambda}.$$

This combine with (3.8) and (3.9), on $\mathbb{R}_+^{n+1}\cap\partial B_1$, we have

$$\begin{aligned}\frac{\partial^2}{\partial r^2}\frac{du_e^\lambda}{d\lambda}=&\lambda\frac{\partial}{\partial r}\frac{d^2u_e^\lambda}{d\lambda^2}-\frac{2s}{p-1}\frac{\partial}{\partial r}\frac{du_e^\lambda}{d\lambda}\\ =&\lambda^2\frac{d^3u_e^\lambda}{d\lambda^3}+\lambda(2-\frac{4s}{p-1})\frac{d^2u_e^\lambda}{d\lambda^2}-\frac{2s}{p-1}(1-\frac{2s}{p-1})\frac{du_e^\lambda}{d\lambda}.\end{aligned}\tag{3.19}$$

Now from (3.17), combine with (3.18) and (3.19), we get

$$\begin{aligned}\frac{\partial^3}{\partial r^3}\frac{du_e^\lambda}{d\lambda}=&\lambda^3\frac{d^4u_e^\lambda}{d\lambda^4}+\lambda^2(3-\frac{6s}{p-1})\frac{d^3u_e^\lambda}{d\lambda^3}-\lambda(1-\frac{2s}{p-1})\frac{6s}{p-1}\frac{d^2u_e^\lambda}{d\lambda^2}\\ &+(1-\frac{2s}{p-1})(1+\frac{2s}{p-1})\frac{2s}{p-1}\frac{du_e^\lambda}{d\lambda}.\end{aligned}\tag{3.20}$$

From (3.13), on $\mathbb{R}_+^{n+1}\cap\partial B_1$, combine with (3.20) then

$$\begin{aligned}\frac{\partial^4}{\partial r^4}u_e^\lambda=&\lambda\frac{\partial^3}{\partial r^3}\frac{du_e^\lambda}{d\lambda}-(3+\frac{2s}{p-1})\frac{\partial^3}{\partial r^3}u_e^\lambda\\ =&\lambda^4\frac{d^4u_e^\lambda}{d\lambda^4}-\lambda^3\frac{8s}{p-1}\frac{d^3u_e^\lambda}{d\lambda^3}+\lambda^2(2+\frac{4s}{p-1})\frac{6s}{p-1}\frac{d^2u_e^\lambda}{d\lambda^2}\\ &-\lambda(1+\frac{2s}{p-1})(1+\frac{s}{p-1})\frac{16s}{p-1}\frac{du_e^\lambda}{d\lambda}\\ &+(3+\frac{2s}{p-1})(2+\frac{2s}{p-1})(1+\frac{2s}{p-1})\frac{2s}{p-1}u_e^\lambda.\end{aligned}$$

In summary, we have that

$$\begin{aligned}\frac{\partial^3}{\partial r^3}u_e^\lambda=&\lambda^3\frac{d^3u_e^\lambda}{d\lambda^3}-\lambda^2\frac{6s}{p-1}\frac{d^2u_e^\lambda}{d\lambda^2}+\lambda(\frac{6s}{p-1}+\frac{12s^2}{(p-1)^2})\frac{du_e^\lambda}{d\lambda}\\ &-(2+\frac{2s}{p-1})(1+\frac{2s}{p-1})\frac{2s}{p-1}u_e^\lambda\end{aligned}$$

and

$$\frac{\partial^2}{\partial r^2}u_e^\lambda = \lambda^2\frac{d^2u_e^\lambda}{d\lambda^2} - \lambda\frac{4s}{p-1}\frac{du_e^\lambda}{d\lambda} + (1+\frac{2s}{p-1})\frac{2s}{p-1}u_e^\lambda$$

$$\frac{\partial u_e^\lambda}{\partial r} = \lambda\frac{du_e^\lambda}{d\lambda} - \frac{2s}{p-1}u_e^\lambda.$$

3.3 On the operator Δ_b^2 and its representation

Note that

$$\Delta_b u = y^{-b}\nabla\cdot(y^b\nabla u) = \quad u_{rr} + \frac{n+b}{r}u_r + \frac{1}{r^2}\theta_1^{-b}\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}u),$$

where $\theta_1 = \frac{y}{r}, r = \sqrt{|x|^2+y^2}$. Set $v = \Delta_b u$ and $\Delta_b^2 u := w$. Then

$$\begin{aligned}
w =&\Delta_b v = v_{rr} + \frac{n+b}{r}v_r + \frac{1}{r^2}\theta_1^{-b}\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}v)\\
=&\partial_{rrrr}u + \frac{2(n+b)}{r}\partial_{rrr}u + \frac{(n+b)(n+b-2)}{r^2}\partial_{rr}u - \frac{(n+b)(n+b-2)}{r^3}\partial_r u\\
&+ r^{-4}\theta_1^{-b}\mathbf{div}_{S^n}(\theta_1^b\nabla(\theta_1^{-b}\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}u))\\
&+ 2r^{-2}\theta_1^{-b}\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}(u_{rr} + \frac{n+b-2}{r}u_r))\\
&- 2(n+b-3)r^{-4}\theta_1^{-b}\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}u).
\end{aligned}$$

On $\mathbb{R}^{n+1}_+ \cap \partial B_1$, we have

$$\begin{aligned}
w =&\underbrace{\partial_{rrrr}u + 2(n+b)\partial_{rrr}u + (n+b)(n+b-2)\partial_{rr}u - (n+b)(n+b-2)\partial_r u}\\
&\underbrace{+\theta_1^{-b}\mathbf{div}_{S^n}(\theta_1^b\nabla(\theta_1^{-b}\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}u))}\\
&\underbrace{+2\theta_1^{-b}\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}(u_{rr} + \frac{n+b-2}{r}u_r))}\\
&\underbrace{-2(n+b-3)\theta_1^{-b}\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}u)}\\
:=&I(u) + J(u) + K(u) + L(u).
\end{aligned}$$

By these notations, we can rewrite the term $\overline{E}_{d2}(u_e^\lambda,1)$ appear in (3.6) as following

$$
\begin{aligned}
&\overline{E}_{d2}(u_e^\lambda,1)\\
&=\int_{\mathbb{R}_+^{n+1}\cap\partial B_1}\theta_1^b(\lambda\frac{dw_e^\lambda}{d\lambda}\frac{du_e^\lambda}{d\lambda}-\lambda w_e^\lambda\frac{d^2u_e^\lambda}{d\lambda^2})-5\theta_1^b w_e^\lambda\frac{du_e^\lambda}{d\lambda}\\
&=\underbrace{\int_{\mathbb{R}_+^{n+1}\cap\partial B_1}\lambda\theta_1^b\frac{d}{d\lambda}I(u_e^\lambda)\frac{du_e^\lambda}{d\lambda}-\lambda\theta_1^b I(u_e^\lambda)\frac{d^2u_e^\lambda}{d\lambda^2}-5\theta_1^b I(u_e^\lambda)\frac{du_e^\lambda}{d\lambda}}\\
&\underbrace{+\int_{\mathbb{R}_+^{n+1}\cap\partial B_1}\lambda\theta_1^b\frac{d}{d\lambda}J(u_e^\lambda)\frac{du_e^\lambda}{d\lambda}-\lambda\theta_1^b J(u_e^\lambda)\frac{d^2u_e^\lambda}{d\lambda^2}-5\theta_1^b J(u_e^\lambda)\frac{du_e^\lambda}{d\lambda}}\\
&\underbrace{+\int_{\mathbb{R}_+^{n+1}\cap\partial B_1}\lambda\theta_1^b\frac{d}{d\lambda}K(u_e^\lambda)\frac{du_e^\lambda}{d\lambda}-\lambda\theta_1^b K(u_e^\lambda)\frac{d^2u_e^\lambda}{d\lambda^2}-5\theta_1^b K(u_e^\lambda)\frac{du_e^\lambda}{d\lambda}}\\
&\underbrace{+\int_{\mathbb{R}_+^{n+1}\cap\partial B_1}\lambda\theta_1^b\frac{d}{d\lambda}L(u_e^\lambda)\frac{du_e^\lambda}{d\lambda}-\lambda\theta_1^b L(u_e^\lambda)\frac{d^2u_e^\lambda}{d\lambda^2}-5\theta_1^b L(u_e^\lambda)\frac{du_e^\lambda}{d\lambda}},
\end{aligned}
\tag{3.21}
$$

we define as

$$
\begin{aligned}
\overline{E}_{d2}(u_e^\lambda,1)&:=\mathcal{I}+\mathcal{J}+\mathcal{K}+\mathcal{L}\\
&:=I_1+I_2+I_3+J_1+J_2+J_3+K_1+K_2+K_3+L_1+L_2+L_3.
\end{aligned}
$$

where $I_1,I_2,I_3,J_1,J_2,J_3,K_1,K_2,K_3,L_1,L_2,L_3$ are corresponding successively to the 12 terms in (3.21). By the conclusions of subsection 2.2, we have

$$
\begin{aligned}
I(u_e^\lambda)&=\partial_{rrrr}u_e^\lambda+2(n+b)\partial_{rrr}u_e^\lambda\\
&\quad+(n+b)(n+b-2)\partial_{rr}u_e^\lambda-(n+b)(n+b-2)\partial_r u_e^\lambda\\
&=\lambda^4\frac{d^4u_e^\lambda}{d\lambda^4}+\lambda^3\big(2(n+b)-\frac{8s}{p-1}\big)\big)\frac{d^3u_e^\lambda}{d\lambda^3}\\
&\quad+\lambda^2\big[\frac{12s}{p-1}(1+\frac{2s}{p-1})-(n+b)\frac{12s}{p-1}+(n+b)(n+b-2)\big]\frac{d^2u_e^\lambda}{d\lambda^2}\\
&\quad+\lambda\big[-\frac{8s}{p-1}(1+\frac{2s}{p-1})(2+\frac{2s}{p-1})+2(n+b)\frac{6s}{p-1}(1+\frac{2s}{p-1})\\
&\quad+(n+b)(n+b-2)(-\frac{4s}{p-1}-1)\big]\frac{du_e^\lambda}{d\lambda}\\
&\quad+\big[(1+\frac{2s}{p-1})(2+\frac{2s}{p-1})(3+\frac{2s}{p-1})\frac{2s}{p-1}\\
&\quad-(n+b)(1+\frac{2s}{p-1})(2+\frac{2s}{p-1})\frac{4s}{p-1}\\
&\quad+(n+b)(n+b-2)(\frac{2s}{p-1}+2)\frac{2s}{p-1}\big]u_e^\lambda.
\end{aligned}
\tag{3.22}
$$

For convenience, we denote that

$$I(u_e^\lambda) = \lambda^4 \frac{d^4 u_e^\lambda}{d\lambda^4} + \lambda^3 \delta_1 \frac{d^3 u_e^\lambda}{d\lambda^3} + \lambda^2 \delta_2 \frac{d^2 u_e^\lambda}{d\lambda^2} + \lambda \delta_3 \frac{du_e^\lambda}{d\lambda} + \delta_4 u_e^\lambda, \tag{3.23}$$

where δ_i are the corresponding coefficients of $\lambda^i \frac{d^i u_e^\lambda}{d\lambda^i}$ appeared in (3.22) for $i = 1, 2, 3, 4$. Now taking the derivative of (3.23) with respect to λ, we get

$$\begin{aligned} \frac{d}{d\lambda} I(u_e^\lambda) =& \lambda^4 \frac{d^5 u_e^\lambda}{d\lambda^5} + \lambda^3 (\delta_1 + 4) \frac{d^4 u_e^\lambda}{d\lambda^4} + \lambda^2 (3\delta_1 + \delta_2) \frac{d^3 u_e^\lambda}{d\lambda^3} \\ &+ \lambda (2\delta_2 + \delta_3) \frac{d^2 u_e^\lambda}{d\lambda^2} + (\delta_3 + \delta_4) \frac{du_e^\lambda}{d\lambda} \end{aligned} \tag{3.24}$$

and

$$\begin{aligned} &\partial_{rr} u_e^\lambda + (n + b - 2) \partial_r u_e^\lambda \\ =& \lambda^2 \frac{d^2 u_e^\lambda}{d\lambda^2} + \lambda (n + b - 2 - \frac{4s}{p-1}) \frac{du_e^\lambda}{d\lambda} + \frac{2s}{p-1} (3 + \frac{2s}{p-1} - n - b) u_e^\lambda \\ :=& \lambda^2 \frac{d^2 u_e^\lambda}{d\lambda^2} + \lambda \alpha \frac{du_e^\lambda}{d\lambda} + \beta u_e^\lambda. \end{aligned} \tag{3.25}$$

Hence,

$$\frac{d}{d\lambda} [\partial_{rr} u_e^\lambda + (n + b - 2) \partial_r u_e^\lambda] = \lambda^2 \frac{d^3 u_e^\lambda}{d\lambda^3} + \lambda (\alpha + 2) \frac{d^2 u_e^\lambda}{d\lambda^2} + (\alpha + \beta) \frac{du_e^\lambda}{d\lambda}, \tag{3.26}$$

here $\alpha = n + b - 2 - \frac{4s}{p-1}$ and $\beta = \frac{2s}{p-1}(3 + \frac{2s}{p-1} - n - b)$.

3.4 The computations of I_1, I_2, I_3 and $\mathcal{I}$

$$
\begin{aligned}
I_1 :=& \int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \lambda\theta_1^b \frac{d}{d\lambda} I(u_e^\lambda)\frac{du_e^\lambda}{d\lambda} \\
=& \int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b\big(\lambda^5\frac{d^5u_e^\lambda}{d\lambda^5} + \lambda^4(4+\delta_1)\frac{d^4u_e^\lambda}{d\lambda^4} + \lambda^3(3\delta_1+\delta_2)\frac{d^3u_e^\lambda}{d\lambda^3} \\
&+ \lambda^2(2\delta_2+\delta_3)\frac{d^2u_e^\lambda}{d\lambda^2} + \lambda(\delta_3+\delta_4)\frac{du_e^\lambda}{d\lambda}\big)\frac{du_e^\lambda}{d\lambda} \\
=& \frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b\big[\lambda^5\frac{d^4u_e^\lambda}{d\lambda^4}\frac{du_e^\lambda}{d\lambda} - \lambda^5\frac{d^3u_e^\lambda}{d\lambda^3}\frac{d^2u_e^\lambda}{d\lambda^2} + (\delta_1-1)\lambda^4\frac{d^3u_e^\lambda}{d\lambda^3}\frac{du_e^\lambda}{d\lambda} \\
&+ (4-\delta_1+\delta_2)\lambda^3\frac{d^2u_e^\lambda}{d\lambda^2}\frac{du_e^\lambda}{d\lambda} + \frac{3\delta_1-\delta_2+\delta_3-12}{2}\lambda^2(\frac{du_e^\lambda}{d\lambda})^2\big] \\
&+ \int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b\big[(12-3\delta_1+\delta_2+\delta_4)\lambda(\frac{du_e^\lambda}{d\lambda})^2 \\
&+ (\delta_1-4-\delta_2)\lambda^3(\frac{d^2u_e^\lambda}{d\lambda^2})^2 + \lambda^5(\frac{d^3u_e^\lambda}{d\lambda^3})^2\big] \\
&+ \int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b(6-\delta_1)\lambda^4\frac{d^3u_e^\lambda}{d\lambda^3}\frac{d^2u_e^\lambda}{d\lambda^2},
\end{aligned}
\tag{3.27}
$$

where $\delta_i (i=1,2,3,4)$ are defined in (3.22) and (3.23). Denote $f = u_e^\lambda, f' := \frac{du_e^\lambda}{d\lambda}$, we have used the following differential identities:

$$
\begin{aligned}
\lambda^5 f''''' f' =& \big[\lambda^5 f'''' f' - \lambda^5 f''' f'' - 5\lambda^4 f''' f' + 20\lambda^3 f'' f' - 30\lambda^2 f' f'\big]' \\
&+ 60\lambda(f')^2 - 20\lambda^3(f'')^2 + \lambda^5(f''')^2 + 10\lambda^4 f''' f'',
\end{aligned}
$$

$$
\lambda^4 f'''' f' = \big[\lambda^4 f''' f' - 4\lambda^3 f'' f' + 6\lambda^2 f' f'\big]' - 12\lambda(f')^2 + 4\lambda^3(f'')^2 - \lambda^4 f''' f'',
$$

$$
\lambda^3 f''' f' = \big[\lambda^3 f'' f' - \frac{3\lambda^2}{2} f' f'\big]' + 3\lambda(f')^2 - \lambda^3(f'')^2,
$$

and

$$
\lambda^2 f'' f' = \big[\frac{\lambda^2}{2} f' f'\big]' - \lambda(f')^2.
$$

$$
\begin{aligned}
I_2 &:= -\lambda \int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b I(u_e^\lambda)\frac{d^2u_e^\lambda}{d\lambda^2}\\
&= -\lambda \int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b\big(\lambda^4\frac{d^4u_e^\lambda}{d\lambda^4}+\lambda^3\delta_1\frac{d^3u_e^\lambda}{d\lambda^3}+\lambda^2\delta_2\frac{d^2u_e^\lambda}{d\lambda^2}+\lambda\delta_3\frac{du_e^\lambda}{d\lambda}+\delta_4u_e^\lambda\big)\frac{d^2u_e^\lambda}{d\lambda^2}\\
&= \frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b\big[-\lambda^5\frac{d^3u_e^\lambda}{d\lambda^3}\frac{d^2u_e^\lambda}{d\lambda^2}-\delta_4\lambda\frac{du_e^\lambda}{d\lambda}u_e^\lambda\big]\\
&\quad+\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b\big[\lambda^5(\frac{d^3u_e^\lambda}{d\lambda^3})^2-\delta_2\lambda^3(\frac{d^2u_e^\lambda}{d\lambda^2})^2+\delta_4\lambda(\frac{du_e^\lambda}{d\lambda})^2\big]\\
&\quad+\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b\big[(5-\delta_1)\lambda^4\frac{d^3u_e^\lambda}{d\lambda^3}\frac{d^2u_e^\lambda}{d\lambda^2}-\delta_3\lambda^2\frac{d^2u_e^\lambda}{d\lambda^2}\frac{du_e^\lambda}{d\lambda}+\delta_4\frac{du_e^\lambda}{d\lambda}u_e^\lambda\big].
\end{aligned}
\tag{3.28}
$$

Here we have used that

$$
-\lambda^5 f''''f'' = \big[-\lambda^5 f'''f''\big]' + 5\lambda^4 f'''f'' + \lambda^5(f''')^2
$$

and

$$
-\lambda f''f = [-\lambda f'f]' + f'f + \lambda(f')^2.
$$

$$
\begin{aligned}
I_3 &:= -5\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b I(u_e^\lambda)\frac{du_e^\lambda}{d\lambda}\\
&= -5\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b\big[\lambda^4\frac{d^4u_e^\lambda}{d\lambda^4}+\lambda^3\delta_1\frac{d^3u_e^\lambda}{d\lambda^3}+\lambda^2\delta_2\frac{d^2u_e^\lambda}{d\lambda^2}+\lambda\delta_3\frac{du_e^\lambda}{d\lambda}+\delta_4u_e^\lambda\big]\frac{du_e^\lambda}{d\lambda}\\
&=\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b\big[-5\frac{d^3u_e^\lambda}{d\lambda^3}\frac{du_e^\lambda}{d\lambda}+(20-5\delta_1)\lambda^3\frac{d^2u_e^\lambda}{d\lambda^2}\frac{du_e^\lambda}{d\lambda}\big]\\
&\quad+\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b\big[(5\delta_1-20)\lambda^3(\frac{d^2u_e^\lambda}{d\lambda^2})^2-5\delta_3\lambda(\frac{du_e^\lambda}{d\lambda})^2\big]\\
&\quad+\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b\big[5\lambda^4\frac{d^3u_e^\lambda}{d\lambda^3}\frac{d^2u_e^\lambda}{d\lambda^2}+(15\delta_1-60-5\delta_2)\frac{d^2u_e^\lambda}{d\lambda^2}\frac{du_e^\lambda}{d\lambda}-5\delta_4\frac{du_e^\lambda}{d\lambda}u_e^\lambda\big].
\end{aligned}
\tag{3.29}
$$

Here we have use that

$$
-\lambda^4 f''''f' = \big[-5\lambda^4 f'''f' + 20\lambda^3 f''f'\big]' - 20\lambda^3(f'')^2 - 60\lambda^2 f''f' + 5\lambda^4 f'''f''
$$

and

$$
-\lambda^3 f'''f' = \big[-\lambda^3 f''f'\big]' + 3\lambda^2 f''f' + \lambda^3(f'')^2.
$$

Now we add up I_1, I_2, I_3 and further integrate by part, we can get the term $\mathcal{I}$.

$$\begin{aligned}
\mathcal{I} :=& I_1 + I_2 + I_3 \\
=& \frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b\big[\lambda^5 \frac{d^4u_e^\lambda}{d\lambda^4}\frac{du_e^\lambda}{d\lambda} - 2\lambda^5\frac{d^3u_e^\lambda}{d\lambda^3}\frac{d^2u_e^\lambda}{d\lambda^2} \\
&+ (\delta_1-6)\lambda^4\frac{d^3u_e^\lambda}{d\lambda^3}\frac{du_e^\lambda}{d\lambda} + (24-6\delta_1+\delta_2)\lambda^3\frac{d^2u_e^\lambda}{d\lambda^2}\frac{du_e^\lambda}{d\lambda} \\
&+ (9\delta_1-3\delta_2-36)\lambda^2\frac{du_e^\lambda}{d\lambda}\frac{du_e^\lambda}{d\lambda} \\
&+ (8-\delta_1)\lambda^4(\frac{d^2u_e^\lambda}{d\lambda^2})^2 - \delta_4\lambda\frac{du_e^\lambda}{d\lambda}u_e^\lambda - 2\delta_4(u_e^\lambda)^2\big] \\
&+ \int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b\Big(2\lambda^5(\frac{d^3u_e^\lambda}{d\lambda^3})^2 + (10\delta_1-2\delta_2-56)\lambda^3(\frac{d^2u_e^\lambda}{d\lambda^2})^2 \\
&+ (-18\delta_1+\delta_2-4\delta_3+2\delta_4+72)\lambda(\frac{du_e^\lambda}{d\lambda})^2\Big).
\end{aligned} \tag{3.30}$$

Since $u_e^\lambda(X) = \lambda^{\frac{2s}{p-1}}u_e(\lambda X)$, we have the following

$$\begin{aligned}
\lambda^4\frac{d^4u_e^\lambda}{d\lambda^4} =& \lambda^{\frac{2s}{p-1}}\big[\frac{2s}{p-1}(\frac{2s}{p-1}-1)(\frac{2s}{p-1}-2)(\frac{2s}{p-1}-3)u_e(\lambda X) \\
&+ \frac{8s}{p-1}(\frac{2s}{p-1}-1)(\frac{2s}{p-1}-2)r\lambda\partial_r u_e(\lambda X) \\
&+ \frac{12s}{p-1}(\frac{2s}{p-1}-1)r^2\lambda^2\partial_{rr}u_e(\lambda X) \\
&+ \frac{8s}{p-1}r^3\lambda^3\partial_{rrr}u_e(\lambda X) + r^4\lambda^4\partial_{rrrr}u_e(\lambda X)\big],
\end{aligned}$$

and

$$\begin{aligned}
\lambda^3\frac{d^3u_e^\lambda}{d\lambda^3} =& \lambda^{\frac{2s}{p-1}}\big[\frac{2s}{p-1}(\frac{2s}{p-1}-1)(\frac{2s}{p-1}-2)u_e(\lambda X) \\
&+ \frac{6s}{p-1}(\frac{2s}{p-1}-1)r\lambda\partial_r u_e(\lambda X) \\
&+ \frac{6s}{p-1}r^2\lambda^2\partial_{rr}u_e(\lambda X) + r^3\lambda^3\partial_{rrr}u_e(\lambda X)\big],
\end{aligned}$$

$$\begin{aligned}
&\lambda^2\frac{d^2u_e^\lambda}{d\lambda^2} \\
&= \lambda^{\frac{2s}{p-1}}\big[\frac{2s}{p-1}(\frac{2s}{p-1}-1)u_e(\lambda X) + \frac{4s}{p-1}r\lambda\partial_r u_e(\lambda X) + r^2\lambda^2\partial_{rr}u_e(\lambda X)\big]
\end{aligned}$$

and

$$\lambda\frac{du_e^\lambda}{d\lambda} = \lambda^{\frac{2s}{p-1}}\big[\frac{2s}{p-1}u_e(\lambda X) + r\lambda\partial_r u_e(\lambda X)\big].$$

Hence, by scaling we have the following derivatives:

$$\begin{aligned}
&\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b\lambda^5\frac{d^4u_e^\lambda}{d\lambda^4}\frac{du_e^\lambda}{d\lambda}\\
&\quad=\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_\lambda}\lambda^{2s\frac{p+1}{p-1}-n-5}y^b\big[\frac{2s}{p-1}(\frac{2s}{p-1}-1)(\frac{2s}{p-1}-2)(\frac{2s}{p-1}-3)u_e\\
&\qquad+\frac{8s}{p-1}(\frac{2s}{p-1}-1)(\frac{2s}{p-1}-2)\lambda\partial_r u_e+\frac{12s}{p-1}(\frac{2s}{p-1}-1)\lambda^2\partial_{rr}u_e\\
&\qquad+\frac{8s}{p-1}\lambda^3\partial_{rrr}u_e+\lambda^4\partial_{rrrr}u_e\big]\big[\frac{2s}{p-1}u_e+r\lambda\partial_r u_e\big];
\end{aligned}$$

$$\begin{aligned}
&\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b\lambda^5\frac{d^3u_e^\lambda}{d\lambda^3}\frac{d^2u_e^\lambda}{d\lambda^2}\\
&\quad=\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_\lambda}\lambda^{2s\frac{p+1}{p-1}-n-5}y^b\big[\frac{2s}{p-1}(\frac{2s}{p-1}-1)(\frac{2s}{p-1}-2)u_e\\
&\qquad+\frac{6s}{p-1}(\frac{2s}{p-1}-1)\lambda\partial_r u_e+\frac{6s}{p-1}\lambda^2\partial_{rr}u_e+\lambda^3\partial_{rrr}u_e\big]\\
&\qquad\big[\frac{2s}{p-1}(\frac{2s}{p-1}-1)u_e+\frac{4s}{p-1}\lambda\partial_r u_e+\lambda^2\partial_{rr}u_e\big];
\end{aligned}\tag{3.31}$$

$$\begin{aligned}
&\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b\lambda^4\frac{d^3u_e^\lambda}{d\lambda^3}\frac{du_e^\lambda}{d\lambda}\\
&\quad=\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_\lambda}\lambda^{2s\frac{p+1}{p-1}-n-5}y^b\big[\frac{2s}{p-1}(\frac{2s}{p-1}-1)(\frac{2s}{p-1}-2)u_e\\
&\qquad+\frac{6s}{p-1}(\frac{2s}{p-1}-1)\lambda\partial_r u_e+\frac{6s}{p-1}\lambda^2\partial_{rr}u_e+\lambda^3\partial_{rrr}u_e\big]\\
&\qquad\big[\frac{2s}{p-1}u_e+\lambda\partial_r u_e\big];
\end{aligned}$$

$$\begin{aligned}
&\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b\lambda^3\frac{d^2u_e^\lambda}{d\lambda^2}\frac{du_e^\lambda}{d\lambda}\\
&\quad=\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_\lambda}\lambda^{2s\frac{p+1}{p-1}-n-5}y^b\big[\frac{2s}{p-1}(\frac{2s}{p-1}-1)u_e\\
&\qquad+\frac{4s}{p-1}\lambda\partial_r u_e+\lambda^2\partial_{rr}u_e\big][\frac{2s}{p-1}u_e+\lambda\partial_r u_e\big]
\end{aligned}$$

and

$$\begin{aligned}
&\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b\lambda^2\frac{du_e^\lambda}{d\lambda}\frac{du_e^\lambda}{d\lambda}\\
&\quad=\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_\lambda}\lambda^{2s\frac{p+1}{p-1}-n-5}y^b\big[\frac{2s}{p-1}u_e+\lambda\partial_r u_e\big]^2.
\end{aligned}$$

Further,

$$\begin{aligned}
&\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b\lambda^4\frac{d^2u_e^\lambda}{d\lambda^2}\frac{d^2u_e^\lambda}{d\lambda^2}\\
&\quad=\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_\lambda}\lambda^{2s\frac{p+1}{p-1}-n-5}y^b\big[\frac{2s}{p-1}(\frac{2s}{p-1}-1)u_e\\
&\quad+\frac{4s}{p-1}\lambda\partial_r u_e+\lambda^2\partial_{rr}u_e\big]^2,\\
&\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b\lambda\frac{du_e^\lambda}{d\lambda}u_e^\lambda\\
&\quad=\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_\lambda}\lambda^{2s\frac{p+1}{p-1}-n-5}y^b\big[\frac{2s}{p-1}u_e+\lambda\partial_r u_e\big]u_e,
\end{aligned}$$

and

$$\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b u_e^\lambda=\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_\lambda}\lambda^{2s\frac{p+1}{p-1}-n-5}y^b u_e^2.$$

3.5 The computations of $J_i, K_i, L_i (i=1,2,3)$ and $\mathcal{J},\mathcal{K},\mathcal{L}$

Firstly,

$$\begin{aligned}
J_1:=&\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\lambda\theta_1^b\frac{d}{d\lambda}J(u_e^\lambda)\frac{du_e^\lambda}{d\lambda}=\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\lambda\theta_1^b J(\frac{du_e^\lambda}{d\lambda})\frac{du_e^\lambda}{d\lambda}\\
=&\lambda\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\mathbf{div}_{S^n}\big(\theta_1^b\nabla_{S^n}(\theta_1^{-b}\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}\frac{du_e^\lambda}{d\lambda}))\big)\frac{du_e^\lambda}{d\lambda}\\
=&-\lambda\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b\nabla_{S^n}\big(\theta_1^{-b}\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}\frac{du_e^\lambda}{d\lambda})\big)\nabla_{S^n}\frac{du_e^\lambda}{d\lambda} \qquad (3.32)\\
=&\lambda\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^{-b}\big[\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}\frac{du_e^\lambda}{d\lambda})\big]^2\\
=&\lambda\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b(\Delta_{S^n}\frac{du_e^\lambda}{d\lambda})^2,
\end{aligned}$$

here we have used integrate by part formula on the unit sphere S^n.

$$
\begin{aligned}
J_2 :=& -\lambda \int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b J(u_e^\lambda)\frac{d^2u_e^\lambda}{d\lambda^2}\\
=& -\lambda \int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \mathbf{div}_{S^n}\big(\theta_1^b\nabla_{S^n}(\theta_1^{-b}\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}u_e^\lambda))\big)\frac{d^2u_e^\lambda}{d\lambda^2}\\
=&\lambda\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b\nabla_{S^n}(\theta_1^{-b}\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}u_e^\lambda)\nabla_{S^n}\frac{d^2u_e^\lambda}{d\lambda^2}\\
=& -\lambda \int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^{-b}\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}u_e^\lambda)\frac{d^2}{d\lambda^2}\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}u_e^\lambda)\\
=&\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} -\lambda\theta_1^{-b}\big[\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}u_e^\lambda)\big]\frac{d}{d\lambda}\big[\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}u_e^\lambda)\big] \qquad (3.33)\\
&+\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^{-b}\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}u_e^\lambda)\cdot\frac{d}{d\lambda}\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}u_e^\lambda)\\
&+\lambda\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^{-b}\big[\frac{d}{d\lambda}\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}u_e^\lambda)\big]^2\\
=&\frac{d}{d\lambda}\Big(\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b\Big(-\frac{1}{2}\lambda\frac{d}{d\lambda}(\Delta_{S^n}u_e^\lambda)^2+\frac{1}{2}(\Delta_{S^n}u_e^\lambda)^2\Big)\Big)\\
&+\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b\lambda(\Delta_{S^n}\frac{du_e^\lambda}{d\lambda})^2,
\end{aligned}
$$

here we denote that $g=\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}u_e^\lambda), g'=\frac{d}{d\lambda}\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}u_e^\lambda)$ and we have used that

$$
-\lambda gg' = \big[-gg'\big]' + gg' + \lambda(g')^2 = \big[-gg'+\frac{1}{2}g^2\big]' + \lambda(g')^2.
$$

Further,

$$
\begin{aligned}
J_3 :=& -5\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b J(u_e^\lambda)\frac{du_e^\lambda}{d\lambda}\\
=& -5\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \mathbf{div}_{S^n}\big(\theta_1^b\nabla_{S^n}(\theta_1^{-b}\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}u))\big)\frac{du_e^\lambda}{d\lambda}\\
=&5\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b\nabla_{S^n}\big(\theta_1^{-b}\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}u_e^\lambda)\big)\nabla_{S^n}\frac{du_e^\lambda}{d\lambda} \qquad (3.34)\\
=& -5\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^{-b}\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}u_e^\lambda)\frac{d}{d\lambda}\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}u_e^\lambda)\\
=& -\frac{5}{2}\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b(\Delta_{S^n}u_e^\lambda)^2.
\end{aligned}
$$

Therefore, combine with (3.32), (3.33) and (3.34), we get that

$$\begin{aligned}
\mathcal{J} :=& J_1 + J_2 + J_3 \\
=& 2\lambda \int_{\mathbb{R}^{n+1}_+ \cap \partial B_1} \theta_1^{-b} \big[\frac{d}{d\lambda} \mathbf{div}_{S^n}(\theta_1^b \nabla_{S^n} u_e^\lambda)\big]^2 \\
& - 4 \int_{\mathbb{R}^{n+1}_+ \cap \partial B_1} \theta_1^{-b} \mathbf{div}_{S^n}(\theta_1^b \nabla_{S^n} u) \frac{d}{d\lambda} \mathbf{div}_{S^n}(\theta_1^b \nabla_{S^n} u) \\
& + \frac{d}{d\lambda} \int_{\mathbb{R}^{n+1}_+ \cap \partial B_1} -\lambda \theta_1^b \big[\mathbf{div}_{S^n}(\theta_1^b \nabla_{S^n} u_e^\lambda)\big] \frac{d}{d\lambda} \big[\mathbf{div}_{S^n}(\theta_1^b \nabla_{S^n} u_e^\lambda)\big] \\
=& 2\lambda \int_{\mathbb{R}^{n+1}_+ \cap \partial B_1} \theta_1^{-b} \big[\frac{d}{d\lambda} \mathbf{div}_{S^n}(\theta_1^b \nabla_{S^n} u_e^\lambda)\big]^2 \\
& - 2\frac{d}{d\lambda} \int_{\mathbb{R}^{n+1}_+ \cap \partial B_1} \theta_1^{-b} \mathbf{div}_{S^n}(\theta_1^b \nabla_{S^n} u_e^\lambda) \mathbf{div}_{S^n}(\theta_1^b \nabla_{S^n} u_e^\lambda) \\
& + \frac{d}{d\lambda} \int_{\mathbb{R}^{n+1}_+ \cap \partial B_1} -\lambda \theta_1^{-b} \big[\mathbf{div}_{S^n}(\theta_1^b \nabla_{S^n} u_e^\lambda)\big] \frac{d}{d\lambda} \big[\mathbf{div}_{S^n}(\theta_1^b \nabla_{S^n} u_e^\lambda)\big] \\
=& \frac{d}{d\lambda} \int_{\mathbb{R}^{n+1}_+ \cap \partial B_1} \theta_1^b \Big(-2(\Delta_{S^n} u_e^\lambda)^2 - \frac{1}{2}\lambda \frac{d}{d\lambda} (\Delta_{S^n} u_e^\lambda)^2 \Big) \\
& + 2\lambda \int_{\mathbb{R}^{n+1}_+ \cap \partial B_1} \theta_1^b (\Delta_{S^n} u_e^\lambda)^2.
\end{aligned} \tag{3.35}$$

Note that

$$\begin{aligned}
& \frac{d}{d\lambda} \int_{\mathbb{R}^{n+1}_+ \cap \partial B_1} \theta_1^b \big[\theta_1^{-b} \mathbf{div}_{S^n}(\theta_1^b \nabla_{S^n} u_e^\lambda)\big]^2 \\
&= \frac{d}{d\lambda} \int_{\mathbb{R}^{n+1}_+ \cap \partial B_\lambda} \lambda^{2s\frac{p+1}{p-1} - n - 5} \big(\lambda^2 \Delta_b u_e - \lambda^2 \partial_{rr} u_e - (n+b)\lambda \partial_r u_e\big)^2,
\end{aligned} \tag{3.36}$$

and

$$\begin{aligned}
& \frac{d}{d\lambda} \int_{\mathbb{R}^{n+1}_+ \cap \partial B_1} \theta_1^b \lambda \frac{d}{d\lambda} \big[\theta_1^{-b} \mathbf{div}_{S^n}(\theta_1^b \nabla_{S^n} u_e^\lambda)\big]^2 \\
&= \frac{d}{d\lambda} \int_{\mathbb{R}^{n+1}_+ \cap \partial B_\lambda} \lambda^{2s\frac{p+1}{p-1} - n - 4} \frac{d}{d\lambda} \big(\lambda^2 \Delta_b u_e - \lambda^2 \partial_{rr} u_e - (n+b)\lambda \partial_r u_e\big)^2.
\end{aligned}$$

Next we compute K_1, K_2, K_3 and $\mathcal{K}$.

$$
\begin{aligned}
&K_1\\
:=&\lambda\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b\frac{d}{d\lambda}K(u_e^\lambda)\frac{du_e^\lambda}{d\lambda}\\
=&2\lambda\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\mathbf{div}_{S^n}\big(\theta_1^b\nabla_{S^n}(\frac{d}{d\lambda}(\partial_{rr}+(n+b-2)\partial_r)u_e^\lambda)\big)\frac{du_e^\lambda}{d\lambda}\\
=&2\lambda\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\mathbf{div}_{S^n}\big(\theta_1^b\nabla_{S^n}(\lambda^3\frac{d^3u_e^\lambda}{d\lambda^3}+\lambda^2(\alpha+2)\frac{d^2u_e^\lambda}{d\lambda^2}+\lambda(\alpha+\beta)\frac{du_e^\lambda}{d\lambda})\big)\frac{du_e^\lambda}{d\lambda}\\
=&-2\lambda\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b\nabla_{S^n}\big(\lambda^3\frac{d^3u_e^\lambda}{d\lambda^3}+\lambda^2(\alpha+2)\frac{d^2u_e^\lambda}{d\lambda^2}+\lambda(\alpha+\beta)\frac{du_e^\lambda}{d\lambda}\big)\nabla_{S^n}\frac{du_e^\lambda}{d\lambda}\\
=&\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}-\lambda^3\theta_1^b\frac{d}{d\lambda}\big(\frac{d}{d\lambda}\nabla_{S^n}u_e^\lambda\big)^2+(2-2\alpha)\lambda^2\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b(\frac{d}{d\lambda}\nabla_{S^n}u_e^\lambda)\\
&\cdot(\frac{d^2}{d\lambda^2}\nabla_{S^n}u_e^\lambda)+2\lambda^3\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b(\frac{d^2}{d\lambda^2}\nabla_{S^n}u_e^\lambda)^2\\
&-(2\alpha+2\beta)\lambda\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b(\frac{d}{d\lambda}\nabla_{S^n}u_e^\lambda)^2.
\end{aligned}
\tag{3.37}
$$

Here we denote that $h=\nabla_{S^n}u_e^\lambda, h'=\frac{d}{d\lambda}\nabla_{S^n}u_e^\lambda$, and have used that

$$
-\lambda^3h'h'''=\big[-\frac{\lambda^3}{2}\frac{d}{d\lambda}(h')^2\big]'+3\lambda^2h'h''+\lambda^3(h'')^2.
$$

Next,

$$
\begin{aligned}
K_2:=&-\lambda\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^bK(u_e^\lambda)\frac{d^2u_e^\lambda}{d\lambda^2}\\
=&-2\lambda\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\mathbf{div}_{S^n}(\theta_1^b\nabla_{S^n}\big(\lambda^2\frac{d^2u_e^\lambda}{d\lambda^2}+\lambda\alpha\frac{du_e^\lambda}{d\lambda}+\beta u_e^\lambda\big))\frac{d^2u_e^\lambda}{d\lambda^2}\\
=&2\lambda\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b\nabla_{S^n}\big(\lambda^2\frac{d^2u_e^\lambda}{d\lambda^2}+\lambda\alpha\frac{du_e^\lambda}{d\lambda}+\beta u_e^\lambda\big)\nabla_{S^n}\frac{d^2u_e^\lambda}{d\lambda^2}\\
=&\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b\big[2\beta\lambda\nabla_{S^n}u_e^\lambda\frac{d}{d\lambda}\nabla_{S^n}u_e^\lambda-\beta(\nabla_{S^n}u_e^\lambda)^2\big]\\
&+2\lambda^3\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b(\frac{d^2}{d\lambda^2}\nabla_{S^n}u_e^\lambda)^2-2\lambda\beta\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b(\frac{d}{d\lambda}\nabla_{S^n}u_e^\lambda)^2\\
&+2\lambda^2\alpha\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b\frac{d}{d\lambda}\nabla_{S^n}u_e^\lambda\frac{d^2}{d\lambda^2}\nabla_{S^n}u_e^\lambda.
\end{aligned}
\tag{3.38}
$$

Here we have used that

$$
2\lambda hh''=[2\lambda hh'-h^2]'-2\lambda(h')^2.
$$

Further,

$$\begin{aligned}
K_3 :=& -5\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b K(u_e^\lambda)\frac{du_e^\lambda}{d\lambda}\\
=& -10\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\mathbf{div}_{S^n}\big(\theta_1^b\nabla_{S^n}(\lambda^2\frac{d^2u_e^\lambda}{d\lambda^2}+\lambda\alpha\frac{du_e^\lambda}{d\lambda}+\beta u_e^\lambda)\big)\frac{du_e^\lambda}{d\lambda}\\
=&10\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b\nabla_{S^n}\big(\lambda^2\frac{d^2u_e^\lambda}{d\lambda^2}+\lambda\alpha\frac{du_e^\lambda}{d\lambda}+\beta u_e^\lambda\big)\nabla_{S^n}\frac{du_e^\lambda}{d\lambda}\\
=&\frac{d}{d\lambda}\big[5\beta\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b\nabla_{S^n}u_e^\lambda\nabla_{S^n}u_e^\lambda\big]+10\lambda\alpha\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b\big(\frac{d}{d\lambda}\nabla_{S^n}u_e^\lambda\big)^2\\
&+10\lambda^2\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b\frac{d}{d\lambda}\nabla_{S^n}u_e^\lambda\frac{d^2}{d\lambda^2}\nabla_{S^n}u_e^\lambda.
\end{aligned}\tag{3.39}$$

Now combine with (3.37), (3.38) and (3.39), we get that

$$\begin{aligned}
\mathcal{K} :=& K_1+K_2+K_3\\
=&\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b\Big[-\lambda^3\frac{d}{d\lambda}(\frac{d}{d\lambda}\nabla_{S^n}u_e^\lambda)^2\\
&+2\beta\lambda\nabla_{S^n}u_e^\lambda\frac{d}{d\lambda}\nabla_{S^n}u_e^\lambda+4\beta(\nabla_{S^n}u_e^\lambda)^2+6\lambda^2(\nabla_{S^n}\frac{du_e^\lambda}{d\lambda})^2\Big]\\
&+4\lambda^3\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b(\frac{d^2}{d\lambda^2}\nabla_{S^n}u_e^\lambda)^2\\
&+(8\alpha-4\beta-12)\lambda\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b(\frac{d}{d\lambda}\nabla_{S^n}u_e^\lambda)^2.
\end{aligned}\tag{3.40}$$

Notice that by scaling we have

$$\begin{aligned}
&\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b(\nabla_{S^n}u_e^\lambda)^2\\
&=\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_\lambda}\lambda^{2s\frac{p+1}{p-1}-n-5}y^b\big[\lambda^2|\nabla u_e|^2-\lambda^2|\partial_r u_e|^2\big].
\end{aligned}\tag{3.41}$$

$$\begin{aligned}
&\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b\lambda\frac{d}{d\lambda}(\nabla_{S^n}u_e^\lambda)^2\\
&=\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_\lambda}\lambda^{2s\frac{p+1}{p-1}-n-4}y^b\frac{d}{d\lambda}\big[\lambda^2|\nabla u_e|^2-\lambda^2|\partial_r u_e|^2\big]
\end{aligned}$$

and

$$\begin{aligned}
&\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_1}\theta_1^b\lambda^3\frac{d}{d\lambda}(\frac{d}{d\lambda}\nabla_{S^n}u_e^\lambda)^2\\
&=\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_\lambda}\lambda^{2s\frac{p+1}{p-1}-n-4}y^b\frac{d}{d\lambda}\big[\frac{2s}{p-1}\lambda\nabla u_e+\lambda^2\nabla\partial_r u_e\big]^2.
\end{aligned}$$

Finally, we compute $\mathcal{L}$.

$$\begin{aligned}
L_1 &:= \int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \lambda\theta_1^b \frac{d}{d\lambda} L(u_e^\lambda)\frac{du_e^\lambda}{d\lambda} \\
&= -2(n+b-3)\lambda \int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \mathbf{div}_{S^n}(\theta_1^b \nabla_{S^n}\frac{du_e^\lambda}{d\lambda})\frac{du_e^\lambda}{d\lambda} \\
&= 2(n+b-3)\lambda \int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b (\nabla_{S^n}\frac{du_e^\lambda}{d\lambda})^2;
\end{aligned}$$

$$\begin{aligned}
L_2 &:= \int_{\mathbb{R}^{n+1}_+\cap\partial B_1} -\lambda\theta_1^b L(u_e^\lambda)\frac{d^2u_e^\lambda}{d\lambda^2} \\
&= 2(n+b-3)\lambda \int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \mathbf{div}_{S^n}(\theta_1^b \nabla_{S^n} u_e^\lambda)\frac{d^2u_e^\lambda}{d\lambda^2} \\
&= -2(n+b-3) \int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \lambda\theta_1^b \nabla_{S^n} u_e^\lambda \frac{d^2}{d\lambda^2}\nabla_{S^n} u_e^\lambda \\
&= -(n+b-3) \int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b \frac{d}{d\lambda}\big[2\lambda\nabla_{S^n}u_e^\lambda \nabla_{S^n}\frac{du_e^\lambda}{d\lambda} - (\nabla_{S^n}u_e^\lambda)^2\big] \\
&+ 2(n+b-3)\lambda \int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b |\nabla_{S^n}\frac{du_e^\lambda}{d\lambda}|^2;
\end{aligned}$$

$$\begin{aligned}
L_3 &:= \int_{\mathbb{R}^{n+1}_+\cap\partial B_1} -5\theta_1^b L(u_e^\lambda)\frac{du_e^\lambda}{d\lambda} \\
&= 10(n+b-3) \int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \mathbf{div}_{S^n}(\theta_1^b \nabla_{S^n} u_e^\lambda)\frac{du_e^\lambda}{d\lambda} \\
&= -10(n+b-3) \int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b \nabla_{S^n} u_e^\lambda \nabla_{S^n}\frac{du_e^\lambda}{d\lambda} \\
&= -5(n+b-3)\frac{d}{d\lambda} \int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b [\nabla_{S^n} u_e^\lambda]^2.
\end{aligned}$$

Hence,

$$\begin{aligned}
\mathcal{L} &:= L_1 + L_2 + L_3 \\
&= -(n+b-3)\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b\big[\lambda\frac{d}{d\lambda}(\nabla_{S^n} u_e^\lambda)^2\big] \\
&\quad - 4(n+b-3)\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b[\nabla_{S^n} u_e^\lambda]^2 \\
&\quad + 4(n+b-4)\lambda\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b|\nabla_{S^n}\frac{du_e^\lambda}{d\lambda}|^2 \\
&= -(n+b-3)\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_\lambda} \lambda^{2s\frac{p+1}{p-1}-n-4} y^b \frac{d}{d\lambda}\big[\lambda^2|\nabla u_e|^2 - \lambda^2|\partial_r u_e|^2\big] \\
&\quad - 4(n+b-3)\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_\lambda} \lambda^{2s\frac{p+1}{p-1}-n-5} y^b\big[\lambda^2|\nabla u_e|^2 - \lambda^2|\partial_r u_e|^2\big].
\end{aligned}$$

By rescaling, we have

$$\begin{aligned}
&\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b\big[\lambda\frac{d}{d\lambda}(\nabla_{S^n} u_e^\lambda)^2\big] \\
&= \frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_\lambda} \lambda^{2s\frac{p+1}{p-1}-n-4} y^b \frac{d}{d\lambda}\big[\lambda^2|\nabla u_e|^2 - \lambda^2|\partial_r u_e|^2\big]; \\
&\frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_1} \theta_1^b[\nabla_{S^n} u_e^\lambda]^2 \\
&= \frac{d}{d\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_\lambda} \lambda^{2s\frac{p+1}{p-1}-n-5} y^b\big[\lambda^2|\nabla u_e|^2 - \lambda^2|\partial_r u_e|^2\big].
\end{aligned} \tag{3.42}$$

3.6 The term $\overline{E}_{d_1}$

Notice that on the boundary ∂B_1,

$$\begin{aligned}
v_e^\lambda &= \Delta_b u_e^\lambda \\
&= \lambda^2\frac{d^2u_e^\lambda}{d\lambda^2} + (n+b-\frac{4s}{p-1})\lambda\frac{du_e^\lambda}{d\lambda} + \frac{2s}{p-1}(1+\frac{2s}{p-1}-n-b)u_e^\lambda + \Delta_\theta u_e^\lambda \\
&:= \lambda^2\frac{d^2u_e^\lambda}{d\lambda^2} + \alpha_0\lambda\frac{du_e^\lambda}{d\lambda} + \beta_0 u_e^\lambda + \Delta_\theta u_e^\lambda.
\end{aligned}$$

Integrate by part, it follows that

$$\int_{\partial B_1} y^b\lambda(\frac{dv_e^\lambda}{d\lambda})^2 = \int_{\partial B_1}\Big(\lambda^5(\frac{d^3u_e^\lambda}{d\lambda^3})^2 + (\alpha_0^2-2\alpha_0-2\beta_0-4)\lambda^3(\frac{d^2u_e^\lambda}{d\lambda^2})^2$$
$$+(-\alpha_0^2+\beta_0^2+2\alpha_0+2\beta_0)\lambda(\frac{du_e^\lambda}{d\lambda})^2\Big)$$
$$+\int_{\partial B_1}\Big(-2\lambda^3(\nabla_\theta\frac{d^2u_e^\lambda}{d\lambda^2})^2+(10-2\beta_0)\lambda(\nabla_\theta\frac{du_e^\lambda}{d\lambda})^2\Big)+\int_{\partial B_1}\lambda(\Delta_\theta\frac{du_e^\lambda}{d\lambda})^2$$
$$+\frac{d}{d\lambda}\Big(\int_{\partial B_1}\sum_{0\le i,j\le 2, i+j\le 2}c_{i,j}^1\lambda^{i+j}\frac{d^iu_e^\lambda}{d\lambda^i}\frac{d^ju_e^\lambda}{d\lambda^j}+\sum_{0\le s,t\le 2, s+t\le 2}c_{s,t}^2\lambda^{s+t}\frac{d^su_e^\lambda}{d\lambda^s}\frac{d^tu_e^\lambda}{d\lambda^t}\Big), \tag{3.43}$$

where $c_{i,j}^i, c_{s,t}^2$ depending on a, b hence on p, n.

Proof of Theorem 2.2. Notice that the equation (3.6), combine with the estimates on $\mathcal{I}, \mathcal{J}, \mathcal{K}, \mathcal{L}$ and (3.43), we obtain Theorem 2.2. □

4 Energy estimates and Blow down analysis

In this section, we do some energy estimates for the solutions of (1.1), which are important when we perform a blow-down analysis in the next section.

4.1 Energy estimates

Lemma 4.1. *Let u be a solution of* (1.1) *and u_e satisfy* (2.4)*, then there exists a positive constant C such that*

$$\int_{\partial\mathbb{R}_+^{n+1}}|u_e|^{p+1}\eta^6+\int_{\mathbb{R}_+^{n+1}}y^b|\nabla\Delta_bu_e|^2\eta^6$$
$$\le C\big[\int_{\mathbb{R}_+^{n+1}}y^b|\Delta_bu_e|^2\eta^4|\nabla\eta|^2+\int_{\mathbb{R}_+^{n+1}}y^b|\nabla u_e|^2\frac{|\Delta_b\eta^6|^2}{\eta^6}+\int_{\mathbb{R}_+^{n+1}}y^bu_e^2\frac{|\nabla\Delta_b\eta^6|^2}{\eta^6}$$
$$+\int_{\mathbb{R}_+^{n+1}}y^b|\nabla u_e|^2\eta^2|\nabla\eta|^4+\int_{\mathbb{R}_+^{n+1}}y^b|\nabla^2u_e|^2\eta^4|\nabla\eta|^2$$
$$+\int_{\mathbb{R}_+^{n+1}}y^b|\nabla u_e|^2\eta^4|\nabla^2\eta|^2\big]. \tag{4.1}$$

Proof. Multiply the equation (2.4) with $y^b u_e \eta^6$, where η is a test function, we get that

$$\begin{aligned}
0 =& \int_{\mathbb{R}^{n+1}_+} y^b u_e \eta^6 \Delta_b^3 u_e = \int_{\mathbb{R}^{n+1}_+} u_e \eta^6 \mathbf{div}(y^b \nabla \Delta_b^2 u_e) \\
=& -\int_{\partial\mathbb{R}^{n+1}_+} u_e \eta^6 \frac{\partial}{\partial y} \Delta_b^2 u_e - \int_{\mathbb{R}^{n+1}_+} y^b \nabla(u_e \eta^6) \nabla \Delta_b^2 u_e \\
=& C_{n,s} \int_{\partial\mathbb{R}^{n+1}_+} |u_e|^{p+1} \eta^6 - \int_{\partial\mathbb{R}^{n+1}_+} y^b \frac{\partial(u_e\eta^6)}{\partial y} \Delta_b^2 u_e + \int_{\mathbb{R}^{n+1}_+} y^b \Delta_b(u_e\eta^6) \Delta_b^2 u_e \\
=& C_{n,s} \int_{\partial\mathbb{R}^{n+1}_+} |u_e|^{p+1} \eta^6 + \int_{\mathbb{R}^{n+1}_+} y^b \Delta_b(u_e\eta^6) \Delta_b^2 u_e \\
=& C_{n,s} \int_{\partial\mathbb{R}^{n+1}_+} |u_e|^{p+1} \eta^6 - \int_{\partial\mathbb{R}^{n+1}_+} \Delta_b(u_e\eta^6) y^b \frac{\partial \Delta_b u_e}{\partial y} \\
& - \int_{\mathbb{R}^{n+1}_+} y^b \nabla(\Delta_b(u_e\eta^6)) \nabla \Delta_b u_e \\
=& C_{n,s} \int_{\partial\mathbb{R}^{n+1}_+} |u_e|^{p+1} \eta^6 - \int_{\mathbb{R}^{n+1}_+} y^b \nabla(\Delta_b(u_e\eta^6)) \nabla \Delta_b u_e.
\end{aligned} \tag{4.2}$$

Hence, we have

$$C_{n,s} \int_{\partial\mathbb{R}^{n+1}_+} |u_e|^{p+1} \eta^6 = \int_{\mathbb{R}^{n+1}_+} y^b \nabla(\Delta_b(u_e\eta^6)) \nabla \Delta_b u_e. \tag{4.3}$$

Since $\Delta_b(\xi\eta) = \eta\Delta_b\xi + \xi\Delta_b\eta + 2\nabla\xi\nabla\eta$, we have

$$\Delta_b(u_e\eta^6) = \eta^6 \Delta_b u_e + u_e \Delta_b \eta^6 + 12\eta^5 \nabla u_e \nabla \eta,$$

therefore,

$$\begin{aligned}
\nabla\Delta_b(u_e\eta^6)\nabla\Delta_b u_e =& 6\eta^5 \Delta_b u_e \nabla\eta \nabla \Delta_b u_e + (\eta)^6 (\nabla \Delta_b u_e)^2 + \Delta_b \eta^6 \nabla u_e \nabla \Delta_b u_e \\
& + u_e \nabla \Delta_b \eta^6 \nabla \Delta_b u_e + 60\eta^4 (\nabla\eta \nabla \Delta_b u_e)(\nabla u_e \nabla \eta) \\
& + 12\eta^5 \sum_{i,j} \partial_{ij} u_e \partial_i \eta \partial_j \Delta_b u_e + 12\eta^5 \sum_{i,j} \partial_i u_e \partial_{ij} \eta \partial_j \Delta_b u_e.
\end{aligned} \tag{4.4}$$

here $\partial_j (j = 1, ..., n, n+1)$ denote the derivatives with respect to $x_1, ..., x_n, y$ respectively. A similar way can be applied to deal with the following term $|\nabla\Delta_b(u_e\eta^3)|^2$. On the other hand, by the stability condition, we have

$$p \int_{\mathbb{R}^n} |u|^{p+1} \eta^6 \le \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{\Big(u(x)\eta^3(x) - u(y)\eta^3(y)\Big)^2}{|x-y|^{n+2s}} = \frac{1}{C_{n,s}} \int_{\mathbb{R}^{n+1}_+} y^b |\nabla \Delta_b (u_e \eta^3)|^2. \tag{4.5}$$

(Here we notice that $u_e(x, 0) = u(x)$, see Theorem 2.1, (2.3))

Combine with (4.3), (4.4) and (4.5), we have

$$\begin{aligned}
&\int_{\mathbb{R}^{n+1}_+} y^b|\nabla\Delta_b u_e|^2\eta^6\\
&\le C\varepsilon\int_{\mathbb{R}^{n+1}_+} y^b(\nabla\Delta_b u_e)^2\eta^6 + C(\varepsilon)\big[\int_{\mathbb{R}^{n+1}_+} y^b|\Delta_b u_e|^2\eta^4|\nabla\eta|^2\\
&\quad+\int_{\mathbb{R}^{n+1}_+} y^b|\nabla u_e|^2(\frac{|\Delta_b\eta^6|^2}{\eta^6}+\eta^4|\nabla^2\eta|^2)\\
&\quad+\int_{\mathbb{R}^{n+1}_+} y^b u_e^2\frac{|\nabla\Delta_b\eta^6|^2}{\eta^6}+\int_{\mathbb{R}^{n+1}_+} y^b|\nabla u_e|^2\eta^2|\nabla\eta|^4+\int_{\mathbb{R}^{n+1}_+} y^b|\nabla^2 u_e|^2\eta^4|\nabla\eta|^2\big],
\end{aligned}$$

we can select ε so small that $C\varepsilon\le\frac{1}{2}$. Combine with (4.3) and (4.4), we obtain our conclusion. □

Corollary 4.1. *Let u be a solution of* (1.1) *and u_e satisfy* (2.4)*, then*

$$\begin{aligned}
&\int_{\partial\mathbb{R}^{n+1}_+\cap B_{R/2}}|u_e|^{p+1}+\int_{\mathbb{R}^{n+1}_+\cap B_{R/2}} y^b(\nabla\Delta_b u_e)^2\\
&\le C\big[R^{-6}\int_{\mathbb{R}^{n+1}_+\cap B_R} y^b u_e^2+R^{-4}\int_{\mathbb{R}^{n+1}_+\cap B_R} y^b|\nabla u_e|^2\\
&+R^{-2}\int_{\mathbb{R}^{n+1}_+\cap B_R} y^b(|\Delta_b u_e|^2+|\nabla^2 u_e|^2)\big].
\end{aligned}$$

Proof. We let $\eta=\xi^m$ where $m>1$ in the estimate (4.1). We have

$$\begin{aligned}
&\int_{\partial\mathbb{R}^{n+1}_+}|u_e|^{p+1}\xi^{6m}+\int_{\mathbb{R}^{n+1}_+} y^b|\nabla\Delta_b u_e|^2\xi^{6m}\\
&\le C[\int_{\mathbb{R}^{n+1}_+} y^b(|\Delta_b u_e|^2+|\nabla^2 u_e|^2)\xi^{6m-2}|\nabla\xi|^2\\
&\quad+\int_{\mathbb{R}^{n+1}_+} y^b|\nabla u_e|^2\xi^{6m-4}(|\nabla^2\xi|^2+|\nabla\xi|^4)+\int_{\mathbb{R}^{n+1}_+} y^b u_e^2\xi^{6m-6}|\nabla^3\xi|^2].
\end{aligned}$$

Let $\xi=1$ in $B_{R/2}$ and $\xi=0$ in B_R^C, satisfying $|\nabla\xi|\le\frac{C}{R}$, then we have the desired estimates. □

Lemma 4.2. *Suppose that u is a solution of* (1.1) *which is stable outside some ball $B_{R_0}\subset\mathbb{R}^n$. For $\eta\in C_c^\infty(\mathbb{R}^n\backslash\overline{B_{R_0}})$ and $x\in\mathbb{R}^n$, define*

$$\rho(x)=\int_{\mathbb{R}^n}\frac{(\eta(x)-\eta(y))^2}{|x-y|^{n+2s}}dy. \tag{4.6}$$

Then

$$\int_{\mathbb{R}^n}|u|^{p+1}\eta^2dx+\int_{\mathbb{R}^n}\int_{\mathbb{R}^n}\frac{|u(x)\eta(x)-u(y)\eta(y)|^2}{|x-y|^{n+2s}}dxdy\le C\int_{\mathbb{R}^n}u^2\rho dx. \tag{4.7}$$

Lemma 4.3. *Let $m > n/2$ and $x \in \mathbb{R}^n$. Set*

$$\rho(x) := \int_{\mathbb{R}^n} \frac{(\eta(x) - \eta(y))^2}{|x-y|^{n+2s}} dy \quad \text{where} \quad \eta(x) = (1+|x|^2)^{-m/2}. \tag{4.8}$$

Then there is a constant $C = C(n,s,m) > 0$ such that

$$C^{-1}(1+|x|^2)^{-n/2-s} \le \rho(x) \le C(1+|x|^2)^{-n/2-s}. \tag{4.9}$$

Corollary 4.2. *Suppose that $m > n/2$, η is given by* (4.8) *and $R > R_0 > 1$. Define*

$$\rho_R(x) = \int_{\mathbb{R}^n} \frac{(\eta_R(x) - \eta_R(y))^2}{|x-y|^{n+2s}} dy, \quad \text{where} \quad \eta_R(x) = \eta(x/R)\psi(x/R) \tag{4.10}$$

for a standard test function ψ that $\psi \in C^\infty(\mathbb{R}^n)$, $0 \le \psi \le 1, \psi = 0$ on B_1 and $\psi = 1$ on $\mathbb{R}^n \setminus B_2$. Then there exists a constant $C > 0$ such that

$$\rho_R(x) \le C\eta^2(x/R)|x|^{-(n+2s)} + R^{-2s}\rho(x/R).$$

Lemma 4.4. *Suppose that u is a solution of* (1.1) *which is stable outside a ball B_{R_0}. Consider ρ_R which is defined in* (4.10) *for $n/2 < m < n/2 + s(p+1)/2$. Then there exists a constant $C > 0$ such that*

$$\int_{\mathbb{R}^n} u^2 \rho_R \le C\Big(\int_{B_{3R_0}} u^2 \rho_R + R^{n-2s\frac{p+1}{p-1}}\Big)$$

for any $R > 3R_0$.

The proofs of Lemma 4.2, Corollary 4.2, Lemma 4.3 and Lemma 4.4 are similar to that of Lemmas $2.1, 2.2, 2.4$ in [6] and we omit the details here.

Lemma 4.5. *Suppose that $p \ne \frac{n+2s}{n-2s}$. Let u be a solution of* (1.1) *which is stable outside a ball B_{R_0} and u_e satisfy* (2.4)*. Then there exists a constant $C > 0$ such that*

$$\int_{B_R} y^b u_e^2 \le CR^{n+6-2s\frac{p+1}{p-1}}, \quad \int_{B_R} y^b |\nabla u_e|^2 \le CR^{n+4-2s\frac{p+1}{p-1}},$$

$$\int_{B_R} y^b (|\nabla^2 u_e|^2 + |\Delta_b u_e|^2) \le CR^{n+2-2s\frac{p+1}{p-1}}.$$

Proof. Recall that the Possion formula for the fractional equation for the case $0 < s < 1$ (see [2]), we can generalize the expression formula to the general case with non-integer positive real number. Therefore,

$$u_e(x,y) = C_{n,s} \int_{\mathbb{R}^n} u(z) \frac{y^{2s}}{(|x-z|^2 + y^2)^{\frac{n+2s}{2}}} dz.$$

Then we have

$$|u_e(x,y)|^2 \le C \int_{\mathbb{R}^n} u^2(z) \frac{y^{2s}}{(|x-z|^2 + y^2)^{\frac{n+2s}{2}}} dz, \tag{4.11}$$

and

$$\partial_y u_e(x,y) = C_{n,s}\int_{\mathbb{R}^n} u(z)\big[\frac{2sy^{2s-1}}{(|x-z|^2+y^2)^{\frac{n+2s}{2}}} - \frac{(n+2s)y^{2s+1}}{(|x-z|^2+y^2)^{\frac{n+2s+2}{2}}}\big]dz,$$

also

$$\partial_{x_j} u_e(x,y) = -C_{n,s}\int_{\mathbb{R}^n} u(z)\frac{(n+2s)(x_j-z_j)y^{2s}}{(|x-z|^2+y^2)^{\frac{n+2s+2}{2}}}dz,$$

for $j = 1, 2, ..., n$. Hence by Hölder's inequality we have

$$|\nabla u_e(x,y)|^2 \le C\int_{\mathbb{R}^n} \frac{u^2(z)y^{2s-2}}{(|x-z|^2+y^2)^{\frac{n+2s}{2}}}dz. \tag{4.12}$$

By a straightforward calculation we have

$$\begin{aligned}\partial_{x_jx_j} u_e(x,y) =& C_{n,s}\int_{\mathbb{R}^n} u(z)\big[\frac{(n+2s)(n+2s+2)(x_j-z_j)^2y^{2s}}{(|x-z|^2+y^2)^{\frac{n+2s+4}{2}}}\\ &- \frac{(n+2s)y^{2s}}{(|x-z|^2+y^2)^{\frac{n+2s+2}{2}}}\big]dz,\end{aligned}$$

$$\begin{aligned}\partial_{x_jy} u_e(x,y) =& C_{n,s}\int_{\mathbb{R}^n} u(z)\big[\frac{(n+2s)(n+2s+2)(x_j-z_j)^2y^{2s+1}}{(|x-z|^2+y^2)^{\frac{n+2s+4}{2}}}\\ &- \frac{2s(n+2s)(x_j-z_j)^2y^{2s-1}}{(|x-z|^2+y^2)^{\frac{n+2s+2}{2}}}\big],\end{aligned}$$

and

$$\begin{aligned}\partial_{yy} u_e(x,y) =& C_{n,s}\int_{\mathbb{R}^n} u(z)\big[\frac{2s(2s-1)y^{2s-2}}{(|x-z|^2+y^2)^{\frac{n+2s}{2}}}\\ &- \frac{(n+2s)(4s+1)y^{2s}}{(|x-z|^2+y^2)^{\frac{n+2s+2}{2}}} + \frac{(n+2s)(n+2s+2)y^{2s+2}}{(|x-z|^2+y^2)^{\frac{n+2s+4}{2}}}\big].\end{aligned}$$

Therefore, we have

$$|\nabla^2 u_e(x,y)| + |\Delta_b u_e(x,y)| \le C\int_{\mathbb{R}^n} |u(z)|\frac{y^{2s-2}}{(|x-z|^2+y^2)^{\frac{n+2s}{2}}}dz.$$

Hence,

$$|\nabla^2 u_e(x,y)|^2 + |\Delta_b u_e(x,y)|^2 \le C\int_{\mathbb{R}^n} u^2(z)\frac{y^{2s-4}}{(|x-z|^2+y^2)^{\frac{n+2s}{2}}}dz. \tag{4.13}$$

Now we turn to estimate the following integration, which provides a unify way to deal with our desired estimates.

Define

$$
\begin{aligned}
A_k :=& \int_{|x|\le R, z\in\mathbb{R}^n} u^2(z)\big[\int_0^R \frac{y^{2k+1}}{(|x-z|^2+y^2)^{\frac{n+2s}{2}}}dy\big]dzdx \\
=& \int_{|x|\le R, z\in\mathbb{R}^n} u^2(z)\big[\int_0^{R^2} \frac{\alpha^k}{(|x-z|^2+\alpha)^{\frac{n+2s}{2}}}d\alpha\big]dzdx \\
\le& \frac{1}{2}\int_{|x|\le R, z\in\mathbb{R}^n} u^2(z)[\int_0^{R^2} \frac{d\alpha}{(|x-z|^2+\alpha)^{\frac{n+2s}{2}-k}}\big]dzdx \\
=& \frac{1}{2}(\frac{n+2s}{2}-k)\int_{|x|\le R, z\in\mathbb{R}^n} u^2(z)\big[(|x-z|^2)^{k-\frac{n+2s}{2}+1} \\
& -(|x-z|^2+R^2)^{k-\frac{n+2s}{2}+1}\big],
\end{aligned}
\tag{4.14}
$$

where $k=0,1,2$. Split the integral to $|x-z|\le 2R$ and $|x-z|>2R$, for the case of $|x-z|\le 2R$, we see that

$$
\begin{aligned}
&\int_{|x|\le R, |x-z|\le 2R} u^2(z)\Big[(|x-z|^2)^{k-\frac{n+2s}{2}+1}-(|x-z|^2+R^2)^{k-\frac{n+2s}{2}+1}\Big] \\
&\le \int_{|x|\le R, |x-z|\le 2R} u^2(z)\Big[(|x-z|^2)^{k-\frac{n+2s}{2}+1}\Big] \\
&\le CR^{2k-2s+2}\int_{|z|\le 3R} u^2(z)dz \\
&\le R^{2k-2s+2}\Big(\int_{B_{3R}} |u|^{p+1}\eta_R^2\Big)^{2/(p+1)}\Big(\int_{B_{3R}} \eta_R^{-4/(p-1)}\Big)^{(p-1)/(p+1)} \\
&\le CR^{2k-2s+2}\Big(\int_{B_{3R}} u^2(z)\rho_R(z)\Big)^{2/(p+1)} \\
&\le CR^{n+2k+2-2s\frac{p+1}{p-1}}.
\end{aligned}
$$

Here we have used Lemma 4.2 and 4.4. For the case of $|x-z|>2R$, by the mean value theorem, we have

$$
\begin{aligned}
&\int_{|x|\le R, |x-z|>2R} u^2(z)\Big[(|x-z|^2)^{k-\frac{n+2s}{2}+1}-(|x-z|^2+R^2)^{k-\frac{n+2s}{2}+1}\Big] \\
&\le R^2\int_{|x|\le R, |x-z|>2R} u^2(z)\Big[(|x-z|^2)^{k-\frac{n+2s}{2}}\Big] \\
&\le CR^{n+2}\int_{|z|\ge R} u_e^2(z)|z|^{2k-n-2s}dz \\
&\le CR^{n+2}\Big[\int_{|z|\ge R}(u_e^{p+1}(z))\Big]^{2/(p+1)}\Big(\int_{|z|\ge R}|z|^{(2k-n-2s)\frac{p+1}{p-1}}\Big)^{(p-1)/(p+1)} \\
&\le CR^{n+2k+2-2s\frac{p+1}{p-1}},
\end{aligned}
$$

here we have used Lemma 4.2. Hence, we obtain that

$$
A_k \le CR^{n+2k+2-2s\frac{p+1}{p-1}}, \tag{4.15}
$$

where $C = C(n, s, p)$ independent of R. Now, combine with (4.11), (4.12) and (4.13), recall that $b = 5 - 2s$, we have

$$\int_{B_R} y^b u_e^2 dxdy \leq A_2, \quad \int_{B_R} y^b |\nabla u_e|^2 dxdy \leq A_1,$$
$$\int_{B_R} y^b \big(|\nabla^2 u_e|^2 + |\Delta_b u_e|^2\big) dxdy \leq A_0.$$

Apply (4.15), we finish our proof. □

Combine Corollary 4.1 and Lemma 4.4, we have the following lemma.

Lemma 4.6. *Let u be a solution of* (1.1) *which is stable outside a ball B_{R_0} and u_e satisfy* (2.4)*. Then there exists a positive constant C such that*

$$\int_{\partial\mathbb{R}^{n+1}_+ \cap B_R} |u_e|^{p+1} + R^{-6} \int_{\mathbb{R}^{n+1}_+ \cap B_R} y^b |u_e|^2 + R^{-4} \int_{\mathbb{R}^{n+1}_+ \cap B_R} y^b |\nabla u_e|^2$$
$$+ R^{-2} \int_{\mathbb{R}^{n+1}_+ \cap B_R} y^b \big(|\Delta_b u_e|^2 + |\nabla^2 u_e|^2\big) + \int_{\mathbb{R}^{n+1}_+ \cap B_R} y^b |\nabla \Delta_b u_e|^2 \leq C R^{n - 2s\frac{p+1}{p-1}}.$$

4.2 Blow down analysis and the proof of Theorem 1.1

The proof of Theorem 1.1. Suppose that u is a solution of (1.1) which is stable outside the ball of radius R_0 and suppose that u_e satisfies (2.4). In the subcritical case, i.e., $1 < p < p_s(n)$, Lemma 4.2 implies that $u \in \dot{H}^s(\mathbb{R}^n) \cap L^{p+1}(\mathbb{R}^n)$. Multiplying (1.1) with u and integrate, we obtain that

$$\int_{\mathbb{R}^n} |u|^{p+1} = \|u\|^2_{\dot{H}^s(\mathbb{R}^n)}. \tag{4.16}$$

Multiplying (1.1) with $u^\lambda(x) = u(\lambda x)$ yields

$$\int_{\mathbb{R}^n} |u|^{p-1}u^\lambda = \int_{\mathbb{R}^n} (-\Delta)^{s/2}u(-\Delta)^{s/2}u^\lambda = \lambda^s \int_{\mathbb{R}^n} w w_\lambda,$$

where $w = (-\Delta)^{s/2}u$. Following the ideas provided in [24, 25] and using the change of variable $z = \sqrt{\lambda}x$, we can get the following Pohozaev identity

$$\begin{aligned} -\frac{n}{p+1}\int_{\mathbb{R}^n} |u|^{p+1} &= \frac{2s-n}{2}\int_{\mathbb{R}^n} |w|^2 + \frac{d}{d\lambda}\int_{\mathbb{R}^n} w^{\sqrt{\lambda}} w^{1/\sqrt{\lambda}} dz\Big|_{\lambda=1} \\ &= \frac{2s-n}{2}\|u\|^2_{\dot{H}^s(\mathbb{R}^n)}. \end{aligned}$$

Hence, we have the following Pohozaev identity

$$\frac{n}{p+1}\int_{\mathbb{R}^n} |u|^{p+1} = \frac{n-2s}{2}\|u\|^2_{\dot{H}^s(\mathbb{R}^n)}.$$

For $p < p_s(n)$, this equality above together with (4.16) proves that $u \equiv 0$. For $p = p_s(n)$, this equality above means that the energy is finite. Further, since $u \in \dot{H}^s(\mathbb{R}^n)$, apply the stability inequality with test function $\psi = u\eta^2(\frac{x}{R})$, and let $R \to +\infty$ (where η is cutoff function), then we get that

$$p\int_{\mathbb{R}^n} |u|^{p+1} \le \|u\|^2_{\dot{H}^s(\mathbb{R}^n)}.$$

This together with (4.16) gives that $u \equiv 0$.

Now we consider the supercritical case, i.e., $p > \frac{n+2s}{n-2s}$, we perform the proof via a few steps.

Step 1. $\lim_{\lambda\to\infty} E(u_e, 0, \lambda) < \infty$.

From Theorem 2.4 we know that E is nondecreasing w.r.t. λ, so we only need to show that $E(u_e, 0, \lambda)$ is bounded. Note that

$$E(u_e, 0, \lambda) \le \frac{1}{\lambda}\int_\lambda^{2\lambda} E(u_e, 0, t)dt \le \frac{1}{\lambda^2}\int_\lambda^{2\lambda}\int_t^{t+\lambda} E(u_e, 0, \gamma)d\gamma dt.$$

From Lemma 4.6, we have that

$$\frac{1}{\lambda^2}\int_{\lambda}^{2\lambda}\int_{t}^{t+\lambda}\gamma^{2s\frac{p+1}{p-1}-n}\big[\int_{\mathbb{R}^{n+1}_{+}\cap B_\gamma}\frac{1}{2}y^b|\nabla\Delta_b u_e|^2dydx - \frac{C_{n,s}}{p+1}\int_{\partial\mathbb{R}^{n+1}_{+}\cap B_\gamma}|u_e|^{p+1}dx\big]d\gamma dt \le C,$$

where $C>0$ is independent of γ.

$$\begin{aligned}
&\frac{1}{\lambda^2}\int_{\lambda}^{2\lambda}\int_{t}^{t+\lambda}\int_{\mathbb{R}^{n+1}_{+}\cap\partial B_\gamma}\gamma^{2s\frac{p+1}{p-1}-n-5}y^b\big[\frac{2s}{p-1}(\frac{2s}{p-1}-1)(\frac{2s}{p-1}-2)u_e\\
&\quad+\frac{6s}{p-1}(\frac{2s}{p-1}-1)\gamma\partial_r u_e+\frac{6s}{p-1}\gamma^2\partial_{rr}u_e+\gamma^3\partial_{rrr}u_e\big]\\
&\qquad\big[\frac{2s}{p-1}(\frac{2s}{p-1}-1)u_e+\frac{4s}{p-1}\gamma\partial_r u_e+\gamma^2\partial_{rr}u_e\big]\\
&\quad\le C\frac{1}{\lambda^2}\int_{\lambda}^{2\lambda}\int_{t}^{t+\lambda}t^{2s\frac{p+1}{p-1}-n-5}\int_{\mathbb{R}^{n+1}_{+}\cap\partial B_\gamma}\\
&\qquad y^b\big[u_e^2+\gamma^2(\partial_r u_e)^2+\gamma^4(\partial_{rr}u_e)^2+\gamma^6(\partial_{rrr}u_e)^2\big]\\
&\quad\le C\frac{1}{\lambda^2}\int_{\lambda}^{2\lambda}t^{2s\frac{p+1}{p-1}-n-5}\int_{\mathbb{R}^{n+1}_{+}\cap B_{3\lambda}}\\
&\qquad y^b\big[u_e^2+\gamma^2(\partial_r u_e)^2+\gamma^4(\partial_{rr}u_e)^2+\gamma^6(\partial_{rrr}u_e)^2\big]\\
&\quad\le C\lambda^{n-2s\frac{p+1}{p-1}+6}\frac{1}{\lambda^2}\int_{\lambda}^{2\lambda}t^{2s\frac{p+1}{p-1}-n-5}dt\\
&\quad\le C
\end{aligned}\tag{4.17}$$

and

$$
\begin{aligned}
&\Big|\frac{1}{\lambda^2}\int_\lambda^{2\lambda}\int_t^{t+\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_\gamma}\gamma^{2s\frac{p+1}{p-1}-n-4}y^b\frac{d}{d\gamma}(\gamma^2\Delta_b u_e-\gamma^2\partial_{rr}u_e-(n+b)\gamma\partial_r u_e)^2\Big|\\
&\le\frac{1}{\lambda^2}\int_\lambda^{2\lambda}t^{2s\frac{p+1}{p-1}-n-5}\int_t^{t+\lambda}\int_{\mathbb{R}^{n+1}_+\cap\partial B_\gamma}y^b\big[2\gamma^2\Delta_b u_e-2\gamma^2\partial_{rr}u_e-(n+b)\gamma\partial_r u_e\big]\\
&\quad\big[\gamma^2\Delta_b u_e-\gamma^2\partial_{rr}u_e-(n+b)\gamma\partial_r u_e\big]\\
&\le\frac{1}{\lambda^2}\int_\lambda^{2\lambda}t^{2s\frac{p+1}{p-1}-n-5}\int_t^{t+\lambda}\int_{\mathbb{R}^{n+1}_+\cap B_{3\lambda}}y^b[2\gamma^2\Delta_b u_e-2\gamma^2\partial_{rr}u_e-(n+b)\gamma\partial_r u_e]\\
&\quad\big[\gamma^2\Delta_b u_e-\gamma^2\partial_{rr}u_e-(n+b)\gamma\partial_r u_e\big]\\
&\le C\frac{1}{\lambda^2}\int_\lambda^{2\lambda}t^{2s\frac{p+1}{p-1}-n-5}\int_t^{t+\lambda}\int_{\mathbb{R}^{n+1}_+\cap B_{3\lambda}}y^b\big[\gamma^4(\Delta_b u_e)^2+\gamma^4(\partial_{rr}u_e)^2+\gamma^2\partial_r u_e\big]\\
&\le C\lambda^{n-2s\frac{p+1}{p-1}+6}\frac{1}{\lambda^2}\int_\lambda^{2\lambda}t^{2s\frac{p+1}{p-1}-n-5}dt\\
&\le C.
\end{aligned}
\tag{4.18}
$$

Integrate by part, by the scaling identity of section 3, for example (3.31), (3.36), (3.41) and (3.42), we can treat the remaining terms by a similar way as the estimates (4.17) and (4.18).

Step 2. There exists a sequence $\lambda_i\to\infty$ such that $(u_e^{\lambda_i})$ converges weakly to a function u_e^∞ in $H^3_{loc}(\mathbb{R}^n;y^b dxdy)$, this is a direct consequence of Lemma 4.6.

Step 3. The function u_e^∞ is homogeneous. Due to the scaling invariance of E (i.e., $E(u_e,0,R\lambda)=E(u_e^\lambda,0,R)$) and the monotonicity formula, for any given $R_2>R_1>0$, we see that

$$
\begin{aligned}
0&=\lim_{i\to\infty}\big(E(u_e,0,R_2\lambda_i)-E(u_e,0,R_1\lambda_i)\big)\\
&=\lim_{i\to\infty}\big(E(u_e^{\lambda_i},0,R_2)-E(u_e^{\lambda_i},0,R_1)\big)\\
&\ge\liminf_{i\to\infty}\int_{(B_{R_2}\setminus B_{R_1})\cap\mathbb{R}^{n+1}_+}y^b r^{2s\frac{p+1}{p-1}-n-6}\big(\frac{2s}{p-1}u_e^{\lambda_i}+r\frac{\partial u_e^{\lambda_i}}{\partial r}\big)^2dydx\\
&\ge\int_{(B_{R_2}\setminus B_{R_1})\cap\mathbb{R}^{n+1}_+}y^b r^{2s\frac{p+1}{p-1}-n-6}\big(\frac{2s}{p-1}u_e^{\infty}+r\frac{\partial u_e^{\infty}}{\partial r}\big)^2dydx.
\end{aligned}
$$

In the last inequality we have used the weak convergence of the sequence $(u_e^{\lambda_i})$ to the function u_e^∞ in $H^3_{loc}(\mathbb{R}^n;y^b dxdy)$. This implies that

$$
\frac{2s}{p-1}\frac{u_e^\infty}{r}+\frac{\partial u_e^\infty}{\partial r}=0\ \text{ a.e. in }\ \mathbb{R}^{n+1}_+.
$$

Therefore, u_e^∞ is homogeneous.

Step 4. $u_e^\infty = 0$**.** This is a direct consequence of Theorem 3.1 in [17].

Step 5. $(u_e^{\lambda_i})$ converges strongly to zero in $H^3(B_R \setminus B_\varepsilon; y^b dxdy)$ and $(u_e^{\lambda_i})$ converges strongly in $L^{p+1}(\partial\mathbb{R}^{n+1}_+ \cap (B_R \setminus B_\varepsilon))$ for all $R > \varepsilon > 0$. These are consequent results of Lemma 4.6 and Theorem 1.5 in [9].

Step 6. $u_e = 0$. Note that

$$\begin{aligned}
\overline{E}(u_e, \lambda) =& \overline{E}(u_e^\lambda, 1)\\
=& \frac{1}{2}\int_{\mathbb{R}^{n+1}_+ \cap B_1} y^b |\nabla \Delta_b u_e^\lambda|^2 dxdy - \frac{C_{n,s}}{p+1}\int_{\partial\mathbb{R}^{n+1}_+ \cap B_1} |u_e^\lambda|^{p+1} dx\\
=& \frac{1}{2}\int_{\mathbb{R}^{n+1}_+ \cap B_\varepsilon} y^b |\nabla \Delta_b u_e^\lambda|^2 dxdy - \frac{C_{n,s}}{p+1}\int_{\partial\mathbb{R}^{n+1}_+ \cap B_\varepsilon} |u_e^\lambda|^{p+1} dx\\
&+ \frac{1}{2}\int_{\mathbb{R}^{n+1}_+ \cap (B_1 \setminus B_\varepsilon)} y^b |\nabla \Delta_b u_e^\lambda|^2 dxdy - \frac{C_{n,s}}{p+1}\int_{\partial\mathbb{R}^{n+1}_+ \cap (B_1 \setminus B_\varepsilon)} |u_e^\lambda|^{p+1} dx\\
=& \varepsilon^{n-2s\frac{p+1}{p-1}} \overline{E}(u_e, \lambda\varepsilon) + \frac{1}{2}\int_{\mathbb{R}^{n+1}_+ \cap (B_1 \setminus B_\varepsilon)} y^b |\nabla \Delta_b u_e^\lambda|^2 dxdy\\
&- \frac{C_{n,s}}{p+1}\int_{\partial\mathbb{R}^{n+1}_+ \cap (B_1 \setminus B_\varepsilon)} |u_e^\lambda|^{p+1} dx.
\end{aligned}$$

Letting $\lambda \to +\infty$ and then $\varepsilon \to 0$, we deduce that $\lim_{\lambda\to+\infty} \overline{E}(u_e, \lambda) = 0$. Using the monotonicity of E,

$$\begin{aligned}
E(u_e, \lambda) \le& \frac{1}{\lambda}\int_\lambda^{2\lambda} E(t)dt \le \sup_{[\lambda,2\lambda]} \overline{E} + C\frac{1}{\lambda}\int_\lambda^{2\lambda} [E - \overline{E}]\\
\le& \sup_{[\lambda,2\lambda]} \overline{E} + C\frac{1}{\lambda}\int_\lambda^{2\lambda} \lambda^{2s\frac{p+1}{p-1}-n-5}\int_{\mathbb{R}^{n+1}_+ \cap \partial B_\lambda} y^b \big[(u_e)^2 + \lambda^2|\nabla u_e|^2\\
&+ \lambda^4(|\Delta_b u_e|^2 + |\nabla^2 u_e|^2) + \lambda^6 |\nabla \Delta_b u_e|^2\big]\\
=& \sup_{[\lambda,2\lambda]} \overline{E} + C\frac{1}{\lambda}\lambda^{2s\frac{p+1}{p-1}-n-5}\int_{\mathbb{R}^{n+1}_+ \cap (B_{2\lambda} \setminus B_\lambda)} y^b \big[(u_e)^2 + \lambda^2|\nabla u_e|^2\\
&+ \lambda^4(|\Delta_b u_e|^2 + |\nabla^2 u_e|^2) + \lambda^6 |\nabla \Delta_b u_e|^2\big]\\
=& \sup_{[\lambda,2\lambda]} \overline{E} + C\frac{1}{\lambda}\int_{\mathbb{R}^{n+1}_+ \cap (B_2 \setminus B_1)} y^b \big[(u_e^\lambda)^2 + |\nabla u_e^\lambda|^2\\
&+ |\Delta_b u_e^\lambda|^2 + |\nabla^2 u_e^\lambda|^2 + |\nabla \Delta_b u_e^\lambda|^2\big]
\end{aligned} \tag{4.19}$$

and so $\lim_{\lambda\to\infty} E(u_e, \lambda) = 0$. Since u is smooth, we also have $E(u_e, 0) = 0$. Since E is monotone, $E \equiv 0$ and so $\overline{u}$ must be homogenous, a contradiction unless $u_e \equiv 0$.

5 Algebraic analysis: The proof of Theorem 2.3

Let $k := \frac{2s}{p-1}$ and $m := n - 2s$. By a direct calculation, we obtain that

$$\begin{aligned} A_1 &= -10k^2 + 10mk - m^2 + 12m + 25, \\ A_2 &= 3k^4 - 6mk^3 + (3m^2 - 12m - 30)k^2 + (12m^2 + 30m)k + 9m^2 + 36m + 27, \\ B_1 &= -6k^2 + 6mk + 12m + 30. \end{aligned} \tag{5.1}$$

Notice that our supercritical condition $p > \frac{n+2s}{n-2s}$ is equivalent to $0 < k < \frac{n-2s}{2} = \frac{m}{2}$. Next, we have the following lemma which yields the sign of A_2 and B_1.

Lemma 5.1. *If $p > \frac{n+2s}{n-2s}$, then $A_2 > 0$ and $B_1 > 0$.*

Proof. From (5.1), we derive that

$$A_2 = 3(k+1)(k+3)(k-(m+1))(k-(m+3)), \tag{5.2}$$

and the roots of $B_1 = 0$ are

$$\frac{1}{2}m - \frac{1}{2}\sqrt{m^2+8m+20}, \quad \frac{1}{2}m + \frac{1}{2}\sqrt{m^2+8m+20}.$$

Recall that $p > \frac{n+2s}{n-2s}$ is equivalent to $0 < k < \frac{m}{2}$, we get the conclusion. □

To show monotonicity formula, we proceed to prove the following inequality. That is, there exist real numbers $c_{i,j}$ and positive real number ϵ such that

$$\begin{aligned} &3\lambda^5(\frac{d^3u^\lambda}{d\lambda^3})^2 + A_1\lambda^3\Big(\frac{d^2u^\lambda}{d\lambda^2}\Big)^2 + A_2\lambda(\frac{du^\lambda}{d\lambda})^2 \\ &\geq \epsilon\lambda(\frac{du^\lambda}{d\lambda})^2 + \frac{d}{d\lambda}\Big(\sum_{0\leq i,j\leq 2} c_{i,j}\lambda^{i+j}\frac{d^iu^\lambda}{d\lambda^i}\frac{d^ju^\lambda}{d\lambda^j}\Big). \end{aligned} \tag{5.3}$$

To deal with the rest of the dimensions, we employ the second idea: we find nonnegative constants d_1, d_2 and constants c_1, c_2 such that we have the following Jordan form decomposition:

$$\begin{aligned} 3\lambda^5(f''')^2 + A_1\lambda^3(f'')^2 + A_2\lambda(f')^2 = 3\lambda(\lambda^2 f''' + c_1\lambda f'')^2 + d_1\lambda(\lambda f'' + c_2 f')^2 \\ + d_2\lambda(f')^2 + \frac{d}{d\lambda}(\sum_{i,j} e_{i,j}\lambda^{i+j}f^{(i)}f^{(j)}), \end{aligned} \tag{5.4}$$

where the unknown constants are to be determined.

Lemma 5.2. *Let $p > \frac{n+2s}{n-2s}$ and A_1 satisfy*

$$A_1 + 12 > 0, \tag{5.5}$$

then there exist nonnegative numbers d_1, d_2, and real numbers $c_1, c_2, e_{i,j}$ such that the differential inequality (5.4) holds.

Proof. Since

$$4\lambda^4 f'''f'' = \frac{d}{d\lambda}(2\lambda^4(f'')^2) - 8\lambda^3(f'')^2$$

and

$$2\lambda^2 f''f' = \frac{d}{d\lambda}(\lambda^2(f')^2) - 2\lambda(f')^2,$$

by comparing the coefficients of $\lambda^3(f'')^2$ and $\lambda(f')^2$, we have that

$$d_1 = A_1 - 3c_1^2 + 12c_1, \quad d_2 = A_2 - (c_2^2 - 2c_2)(A_1 - 3c_1^2 + 12c_1).$$

In particular,

$$\max_{c_1} d_1(c_1) = A_1 + 12 \text{ and the critical point is } c_1 = 2.$$

Since $A_2 > 0$, we select that $c_1 = 2, c_2 = 0$. Hence, in this case, by a direct calculation we see that $d_1 = A_1 + 12 > 0$. Then we get the conclusion. □

We conclude from Lemma 5.2 that if $A_1 + 12 > 0$ then (5.3) holds. This implies that when $m < 6 + \sqrt{73}, p > \frac{n+2s}{n-2s}$ or $m \geq 6 + \sqrt{73}$ and

$$\frac{n+2s}{n-2s} < p < \frac{5m + 20s - \sqrt{15m^2 + 120m + 370}}{5m - \sqrt{15m^2 + 120m + 370}}, \tag{5.6}$$

then (5.3) holds.

Let

$$p_m(n) := \begin{cases} +\infty & \text{if } n < 2s + 6 + \sqrt{73}, \\ \frac{5n+10s-\sqrt{15(n-2s)^2+120(n-2s)+370}}{5n-10s-\sqrt{15(n-2s)^2+120(n-2s)+370}} & \text{if } n \geq 2s + 6 + \sqrt{73}. \end{cases} \tag{5.7}$$

Combining all the lemmas of this section, we obtain the Theorem 2.3.

Now we proceed to prove Theorem 2.4. From Corollary 1.1 of [22], we know that if $n > 2s, s > 0, p > \frac{n+2s}{n-2s}$, then there exists $n_0(s)$, where $\frac{1}{\sqrt{n}} < a_{n,s} < \frac{1}{2}\frac{n-2s}{\sqrt{n}} + \frac{1}{\sqrt{n}}$, such that the inequality (1.4) always holds whenever $n \leq n_0(s)$; while when $n > n_0(s)$, then the inequality (1.4) is true if and only if

$$p < p_2 := \frac{n + 2s - 2 - 2a_{n,s}\sqrt{n}}{n - 2s - 2 - 2a_{n,s}\sqrt{n}},$$

where $n_0(s)$ is in fact the largest n satisfying $n - 2s - 2 - 2a_{n,s}\sqrt{n} \leq 0$. In particular, $\frac{n+2s}{n-2s} < \frac{n+2s-4}{n-2s-4} < p_2 < +\infty$. Therefore, we introduce

$$p_c(n) := \begin{cases} +\infty & \text{if} \quad n \leq n_0(s), \\ \frac{n+2s-2-2a_{n,s}\sqrt{n}}{n-2s-2-2a_{n,s}\sqrt{n}} & \text{if} \quad n > n_0(s). \end{cases} \tag{5.8}$$

From [23], we use the sharp estimate $n_0(s) < 2s + 8.998$ for $2 < s < 3$, then

$$n_0(s) \leq 2s + 8.998 < 2s + 6 + \sqrt{73} \simeq 2s + 14.544. \tag{5.9}$$

On the other hand, via the sharp estimate $a_{n,s} < 1$ from [23]

$$\frac{5n + 10s - \sqrt{15(n-2s)^2 + 120(n-2s) + 370}}{5n - 10s - \sqrt{15(n-2s)^2 + 120(n-2s) + 370}} > \frac{n + 2s - 2 - 2a_{n,s}\sqrt{n}}{n - 2s - 2 - 2a_{n,s}\sqrt{n}} \tag{5.10}$$

provided that $s \in (2, 3)$ and that

$$225m^4 - 720m^3 - 17244m^2 - 29088m + 7236 > 0, \text{ where } m = n - 2s. \tag{5.11}$$

The (5.11) holds whenever $m > 11.12$, that is $n > 2s + 11.12$. This combine with (5.9) we obtain that $p_c(n) < p_m(n)$. Therefore we get Theorem 2.4. □

References

[1] L. Caffarelli, B. Gidas, J. Spruck, Asymptotic symmetry and local behavior of semilinear elliptic equations with critical sobolev growth, *Comm. Pure Appl. Math.*, **42**, no. 3, 271-297, 1989.

[2] L. Caffarelli and L. Silvestre, An extension problem related to the fractional laplacian, *Comm. Partial Differential Equations*, **32**, no. 7-9, 1245-1260, 2007

[3] J. Case, Sun-Yung Alice Chang, On fractional GJMS operators, preprint http://arxiv.org/abs/1406.1846.

[4] Sun-Yung Alice Chang and Maria del Mar Gonzalez, Fractional laplacian in conformal geometry, *Advances in Mathematic,* **226**, no. 2, 1410-1432, 2011.

[5] W. Chen, C. Li, B. Ou, Classification of solutions for an integral equation, *Comm. Pure Appl. Math.*, **59**(2006), no. 3, 330-343.

[6] J. Davila, L. Dupaigne, J. Wei, on the fractional Lane-Emden equation, Trans. Amer. Math. Soc. accepted for publication.

[7] J. Davila, L. Dupaigne, and K. Wang, J. Wei, A monotonicity formula and a liouville-type Theorem for a fourth order supercritical problem, *Advances in Mathematicas*, **258**, 240-285, 2014.

[8] Arpsad Elbert and Andrea Laforgia, On some properties of the gamma function, Proc. Amer. Math. Soc. 128 (2000), 2667-2673.

[9] B. Fabes, E. Kenig and P. Serapioni, The local regularity of solutions of degenerate elliptic equations, *Comm. Partial Differential Equations,* **7**, no. 1, 77-116, 1982.

[10] A. Farina, On the classification of soultions of the Lane-Emden equation on unbounded domains of $\mathbb{R}^N$, *J. Math. Pures Appl.,* **87**(9), no. 5, 537-561, 2007.

[11] M. Fall, Semilinera elliptic equations for the fractional Laplacian with Hardy potential, preprint. http://arxiv.org/pdf/1109.5530v4.pdf

[12] R. Fowler, The solution of Emden's and similar differential equations, *Monthly Notices Roy. Astronom. Soc.*, **91**(1930), 63-91.

[13] R. Fowler, Further studies of Emden's and similar differential equations, *Quarterly J. Math. Oxford Ser.* **2**(1931), 259-288.

[14] B. Gidas, J. Spruck, Global and local behavior of positive solutions of nonlinear elliptic equations, *Comm. Pure Appl. Math.***34**(1981), 525-598.

[15] B. Gidas, J. Spruck, A priori bounds for positive solutions of nonlinear elliptic equations, *Comm. Partial Differential Equations*, **6**(1981), 883-901.

[16] F. Gazzola, H. C. Grunau, Radial entire solutions of supercritical biharmonic equations, *Math. Annal.,* **334**, 905-936, 2006.

[17] M. Fazly, J. Wei, on finite morse index solutions of higher order fractional Lane-Emden equations, American Journal of Mathematics accepted for publication.

[18] D. D. Joseph, T. S. Lundgren, Quasilinear Dirichlet problems driven by positive sources, *Arch. Rational Mech. Anal.,* **49**, 241-269, 1972/73.

[19] Y. Li, Remark on some conformally invariant integral equations: the method of moving spheres, *J. Eur. Math. Soc. (JEMS)* **6**(2004), no. 2, 153-180.

[20] C. S. Lin, A classification of soluitions of a conformally invariant fourth order equation in $\mathbb{R}^N$, *Comment. Math. Helv.,* **73**, 206-231, 1998.

[21] S. Luo, J. Wei and W. Zou, On the Triharmonic Lane-Emden Equation, preprint 2016. arXiv:1607.04719

[22] S. Luo, J. Wei and W. Zou, On a transcendental equation involving quotients of Gamma function, arXiv:1606.06706, Proc. AMS to appear.

[23] S. Luo, J. Wei and W. Zou, On the general Joseph-Lundgren exponent, preprint.

[24] Ros-Oton, J. Serra, Local integration by parts and Pohozaev identities for higher order fractional Laplacians, *Discrete Contin. Dyn. Syst.,* **35**, no. 5, 2131-2150, 2015.

[25] Ros-Oton, J. Serra, The Pohozaev identity for the fractional Laplacian, *Arch. Ration. Mech. Anal.,* **213**, no. 2, 587-628, 2014.

[26] J. Wei, X. Xu, Classification of solutions of higher order conformally invariant equations, *Math. Ann.,* **313**, no. 2, 207-228, 1999.

[27] R. Yang, On higer order extension for the fractional laplacian, preprint, http://arxiv.org/pdf/1302.4413v1.pdf